

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



M

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/66, 1/68, 1/70, C12P 19/34, C12N 15/00, 15/64, 15/85, C07H 21/04		A1	(11) International Publication Number: WO 99/61658
			(43) International Publication Date: 2 December 1999 (02.12.99)
(21) International Application Number: PCT/US99/11629 (22) International Filing Date: 26 May 1999 (26.05.99) (30) Priority Data: 09/085,148 26 May 1998 (26.05.98) US 60/124,090 12 March 1999 (12.03.99) US (71) Applicant: VIROLOGIC, INC. [US/US]; 270 East Grand Avenue, South San Francisco, CA 94080 (US). (72) Inventor: WHITCOMB, Jeanette; 633 Highland Avenue #4, San Mateo, CA 94401 (US). (74) Agent: WHITE, John, P.; Cooper & Dunham LLP, 1185 Avenue of the Americas, New York, NY 10036 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KR, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: MEANS AND METHODS FOR MONITORING NON-NUCLEOSIDE REVERSE TRANSCRIPTASE INHIBITOR ANTIRETROVIRAL THERAPY			
(57) Abstract This invention relates to antiviral drug susceptibility and resistance tests to be used in identifying effective drug regimens for the treatment of human immunodeficiency virus (HIV) infection and acquired immunodeficiency syndrome (AIDS) and further relates to the means and methods of monitoring the clinical progression of HIV infection and its response to antiretroviral therapy, particularly non-nucleoside reverse transcriptase inhibitor therapy using phenotypic susceptibility assays or genotypic assays.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**MEANS AND METHODS FOR MONITORING NON-NUCLEOSIDE REVERSE
TRANSCRIPTASE INHIBITOR ANTIRETROVIRAL THERAPY**

This application is a continuation-in-part of U.S. Serial No. 09/085,148, filed May 26, 1998 and claims the benefit
5 of U.S. Provisional Application No. 60/124,090, filed March 12, 1999, the contents of which are hereby incorporated by reference into this application.

Throughout this application, various references are
10 referred to within parenthesis. Disclosures of these publications in their entireties are hereby incorporated by reference into this application to more fully describe the state of the art to which this invention pertains.

15 **Technical Field**

This invention relates to antiretroviral drug susceptibility and resistance tests to be used in identifying effective drug regimens for the treatment of human immunodeficiency virus (HIV) infection and acquired
20 immunodeficiency syndrome (AIDS). The invention further relates to the means and methods of monitoring the clinical progression of HIV infection and its response to antiretroviral therapy using phenotypic or genotypic susceptibility assays. The invention also relates to novel
25 vectors, host cells and compositions for carrying out phenotypic susceptibility tests. The invention further relates to the use of various genotypic methodologies to identify patients whose infection has become resistant to a particular antiretroviral drug regimen. This invention
30 also relates to the screening of candidate antiretroviral drugs for their capacity to inhibit viruses, selected viral sequences and/or viral proteins. More particularly, this invention relates to the determination of non-nucleoside reverse transcriptase inhibitor resistance using phenotypic
35 susceptibility tests and/or genotypic tests.

Background of the Invention

HIV infection is characterized by high rates of viral turnover throughout the disease process, eventually leading to CD4 depletion and disease progression. Wei X, Ghosh SK, Taylor ME, et al. (1995) *Nature* 343, 117-122 and Ho DD, Naumann AU, Perelson AS, et al. (1995) *Nature* 373, 123-126. The aim of antiretroviral therapy is to achieve substantial and prolonged suppression of viral replication. Achieving sustained viral control is likely to involve the use of sequential therapies, generally each therapy comprising combinations of three or more antiretroviral drugs. Choice of initial and subsequent therapy should, therefore, be made on a rational basis, with knowledge of resistance and cross-resistance patterns being vital to guiding those decisions. The primary rationale of combination therapy relates to synergistic or additive activity to achieve greater inhibition of viral replication. The tolerability of drug regimens will remain critical, however, as therapy will need to be maintained over many years.

In an untreated patient, some 10^{10} new viral particles are produced per day. Coupled with the failure of HIV reverse transcriptase (RT) to correct transcription errors by exonucleolytic proofreading, this high level of viral turnover results in 10^4 to 10^5 mutations per day at each position in the HIV genome. The result is the rapid establishment of extensive genotypic variation. While some template positions or base pair substitutions may be more error prone (Mansky LM, Temin HM (1995) *J Virol* 69, 5087-5094) (Schinazi RF, Lloyd RM, Ramanathan CS, et al. (1994) *Antimicrob Agents Chemother* 38, 268-274), mathematical modeling suggests that, at every possible single point, mutation may occur up to 10,000 times per day in infected individuals.

For antiretroviral drug resistance to occur, the target enzyme must be modified while preserving its function in the presence of the inhibitor. Point mutations leading to

an amino acid substitution may result in change in shape, size or charge of the active site, substrate binding site or surrounding regions of the enzyme. Mutants resistant to antiretroviral agents have been detected at low levels before the initiation of therapy. (Mohri H, Singh MK, Ching WTW, et al. (1993) *Proc Natl Acad Sci USA* 90, 25-29) (Nájera I, Richman DD, Olivares I, et al. (1994) *AIDS Res Hum Retroviruses* 10, 1479-1488) (Nájera I, Holguin A, Quiñones-Mateu E, et al. (1995) *J Virol* 69, 23-31). However, these mutant strains represent only a small proportion of the total viral load and may have a replication or competitive disadvantage compared with wild-type virus. (Coffin JM (1995) *Science* 267, 483-489). The selective pressure of antiretroviral therapy provides these drug-resistant mutants with a competitive advantage and thus they come to represent the dominant quasispecies (Frost SDW, McLean AR (1994) *AIDS* 8, 323-332) (Kellam P, Boucher CAB, Tijnagal JMGH (1994) *J Gen Virol* 75, 341-351) ultimately leading to drug resistance and virologic failure in the patient.

Non-nucleoside Reverse Transcriptase Inhibitors

Non-nucleoside reverse transcriptase inhibitors (NNRTIs) are a chemically diverse group of compounds which are potent inhibitors of HIV-1 RT in vitro. These compounds include pyridinone derivatives, bis(heteroaryl)piperazines (BHAPs) such as delavirdine and atevirdine, the dipyridodiazepinone nevirapine, the thymine derivative groups TSAO and HEPT, an α -anilino phenylacetamides (α -APA) compound loviride, and the quinoxaline-class inhibitors such as (HBY-097), the benzodiazepin-one and -thione (TIBO) compounds and the pyridinone derivatives (L-697,661). For overviews see (DeClercq E. (1996) *Rev Med Virol* 6, 97-117) (Emeni EA (1996) *Antiviral Drug Resistance*, ed. DD Richman, John Wiley & Sons, Ltd. High-level resistance to individual compounds appears to develop rapidly, often within a few weeks of initiating monotherapy, frequently involving only single-point mutations and in many cases

- leading to considerable cross-resistance to other NNRTIs. Most mutations reported occur in the codon groups 100-108 and 181-190 which encode for the two β -sheets adjacent to the catalytic site of the RT enzyme (Kohlstaedt LA, Wang J, Friedman JM, et al. (1992) *Science* 256, 1783-90) The NNRTI binding pocket, as it has been described, is a hydrophobic non-substrate binding region of RT where these agents directly interact with RT. They inhibit activity by interfering with mobility of the 'thumb' subdomain, or disrupting the orientation of conserved aspartic acid side chains essential for catalytic activity (D'Aquila RT. (1994) *Clin Lab Med* 14, 393-423) (Arnold E., Ding J., Hughes SH, et al. (1995) *Curr Opin Struct Biol* 5, 27-38).
- 15 Mutations conferring reduced susceptibility to nevirapine have been described at codons 98, 100, 103, 106, 108, 181, 188 and 190 (Richman DD, Havlir D, Corbeil J. (1994) *J Virol* 68, 1660-1666). The most frequently selected variant during nevirapine monotherapy is a Tyr¹⁸¹Cys (Y181C) mutation which results in a 100-fold reduction in sensitivity to this agent, with reduced susceptibility to the pyridinone derivatives L-696,229 and L-697,661 (Arnold, Ibid). TSAO also has limited activity in the presence of the 181 mutation, but maintains activity in the presence of mutations at codons 100 and 103 and in vitro selects for a unique mutation, GLU¹³⁸Lys (E138K), in the region where it most closely interacts with RT (Richman, DD, Ibid) (Richman DD, Shih C-K, Lowy I, et al. (1991) *Proc Natl Acad Sci USA* 88, 11241-11245).
- 30 Resistance to zidovudine when used as monotherapy develops in most patients by week 24. It has been mapped to a range of codons 100-110; 181-190), most commonly codon 103 (Staszewski S, Miller V, Kober A, et al. (1996) *Antiviral Ther* 1, 42-50. During combination therapy using zidovudine with zidovudine or zidovudine plus lamivudine, variants at codons 98 and 103 were the most frequent mutations detected

at 24 weeks (Staszewski S, Miller V, Rehmet S, et al. (1996) *AIDS* 10, F1-7).

Although the 101, 103 and 181 mutations also confer cross-
5 resistance to BHAPs, (Balzarini J, Karlsson A, Pérez-Pérez M-J, et al. (1992) *Virology* 192, 246-253) the characteristic P236L substitution selected for by these agents *in vitro* appears to sensitize RT to some other NNRTIs, reducing the IC50 for nevirapine, for example, 7-
10 to 10-fold, without influencing sensitivity to nucleoside analogues (Staszewski S., *Ibid*). This mutation at codon 236 has not been observed in clinical isolates during atevirdine therapy, although other resistance-conferring mutations at codons 103 and 181 have been reported during
15 monotherapy as well as at codons 101, 188, 233 and 238 during combination therapy with zidovudine.

While HBY-097 may initially select for a mutation at codons 190 *in vitro*, further passage consistently selects for
20 mutations at RT codon 74 and 75, with some mutant viruses showing decreased sensitivity to didanosine and stavudine, but not zidovudine (Kleim J-P, Rösner M, Winkler I, et al. (1995) *J Acquir Immune Defic Syndr* 10 Suppl 3, 2).

25 Mutation at codon 181 has been reported to antagonize zidovudine resistance due to the typical 41 and 215 codon mutations, (Zhang D, Caliendo AM, Eron JJ, et al. (1994) *Antimicrob Agents Chemother* 38, 282-287) suggesting that combination therapy with some NNRTIs and zidovudine may be
30 feasible. Although an HIV mutant with triple resistance to zidovudine, didanosine and nevirapine has been described *in vitro*, (Larder BA, Kellam P, Kemp SD (1993) *Nature* 365, 451-453) treatment with this triple combination does provide superior immunological and virological responses to
35 treatment with zidovudine plus didanosine alone over a 48-week period in patients with CD4 cell counts <350/mm³.

Combination therapy with zidovudine and the pyridinone derivative L-697,661 prevents the appearance of the codon 181 mutation typically selected during monotherapy with this NNRTI, delaying the appearance of high-level resistance to this compound. Changes in susceptibility to zidovudine were not examined in this study. (Staszewski S, Massari FE, Kober A, et al. (1995) *J Infect Dis* 171, 1159-1165). Concomitant or alternating zidovudine therapy does not delay the appearance of resistance during nevirapine therapy; (Richman DD, Ibid) (Nunberg JH, Schleif WA, Boots EJ, et al. (1990) *J Virol* 65, 4887-4892) (DeJong MD, Loewenthal M, Boucher CAB, et al. (1994) *J Infect Dis* 169, 1346-1350) (Cheeseman SH, Havlir D, McLaughlin MM, et al. (1995) *J Acquir Immune Defic Syndr* 8, 141-151) however, the 181 mutant is not being observed during combination, the most common change being at codon 190 (Richman DD, Ibid). This suggests that the codon 181 mutation which is antagonistic to zidovudine resistance in vitro is not compatible, or not preferred in vivo, selection favoring other mutations which allow for reduced susceptibility to this NNRTI concomitant with zidovudine resistance.

The rapid development of reduced susceptibility to the NNRTIs suggests limited utility of these agents, particularly as monotherapies, and has led to the modification of these molecules in an attempt to delay the appearance of drug-resistant virus. A 'second generation' NNRTI, the pyridinone derivative L-702,019, demonstrated only a 3-fold change in IC_{50} between wild-type and codon 181 mutant HIV-1, and required multiple mutations to engender high-level resistance (Goldman ME, O'Brien JA, Ruffing TL, et al. (1993) *Antimicrob Agents Chemother* 37, 947-949).

It is an object of this invention to provide a drug susceptibility and resistance test capable of showing whether a viral population in a patient is resistant to a given prescribed drug. Another object of this invention is to provide a test that will enable the physician to

substitute one or more drugs in a therapeutic regimen for a patient that has become resistant to a given drug or drugs after a course of therapy. Yet another object of this invention is to provide a test that will enable
5 selection of an effective drug regimen for the treatment of HIV infections and/or AIDS. Yet another object of this invention is to provide the means for identifying the drugs to which a patient has become resistant, in particular identifying resistance to non-nucleoside reverse
10 transcriptase inhibitors. Still another object of this invention is to provide a test and methods for evaluating the biological effectiveness of candidate drug compounds which act on specific viruses, viral genes and/or viral proteins particularly with respect to viral drug resistance
15 associated with non-nucleoside reverse transcriptase inhibitors. It is also an object of this invention to provide the means and compositions for evaluating HIV antiretroviral drug resistance and susceptibility. This and other objects of this invention will be apparent from
20 the specification as a whole.

Summary of the Invention

The present invention relates to methods of monitoring, using phenotypic and genotypic methods, the clinical
25 progression of human immunodeficiency virus infection and its response to antiviral therapy. The invention is also based, in part, on the discovery that genetic changes in HIV reverse transcriptase (RT) which confer resistance to antiretroviral therapy may be rapidly determined directly
30 from patient plasma HIV RNA using phenotypic or genotypic methods. The methods utilize polymerase chain reaction (PCR) based assays. Alternatively, methods evaluating viral nucleic acid of viral protein in the absence of an amplification step could utilize the teaching of this
35 invention to monitor and/or modify antiretroviral therapy. This invention is based in part on the discovery of a mutation at codon 225 either alone or in combination with a mutation at codon 103 of HIV reverse transcriptase in

non-nucleoside reverse transcriptase inhibitor (efavirenz) treated patient(s) in which the presence of the mutations correlate with an increase in delavirdine susceptibility and little or no change in nevirapine susceptibility. The
5 mutations were found in plasma HIV RNA after a period of time following initiation of therapy. The development of the mutant at codon 225 in addition to the mutation at codon 103 in HIV RT was found to be an indicator of the development of resistance and ultimately of immunological
10 decline. This invention is based in part on the discovery of a mutation at codon 236 of RT was discovered to occur in non-nucleoside reverse transcriptase inhibitor (NNRTI) treated patients in which the presence of the mutation correlates with decreased susceptibility to delavirdine and
15 no reduction in nevirapine susceptibility. The development of the codon 190 and 103 and/or 101 mutations in HIV RT was found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with virologic failure and subsequent
20 immunological decline. This invention is based in part on the discovery of a mutation at codon 190 either alone or in combination with a mutation at codon 190 either alone or in combination with a mutation at codon 103 and/or 101 of HIV reverse transcriptase in non-nucleoside reverse
25 transcriptase inhibitor (efavirenz) treated patient(s) in which the presence of the mutations correlate with an increase in delavirdine susceptibility and a decrease in nevirapine susceptibility. The mutations were found in plasma HIV RNA after a period of time following initiation
30 of NNRTI therapy. The development of the codon 236 and 103 and/or 181 mutations in HIV RT was found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline.
35
This invention is based in part on the discovery of a mutation at codon 230 either alone or in combination with a mutation at codon 181 of HIV reverse transcriptase in

non-nucleoside reverse transcriptase inhibitor (nevirapine) treated patient(s) in which the presence of the mutations correlate with a significant decrease in both delavirdine and nevirapine susceptibility. The mutations were found in
5 plasma HIV RNA after a period of time following initiation of NNRTI therapy. The development of the codon 230 and 181 mutations in HIV RT were found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with
10 virologic failure and subsequent immunological decline. This invention is based in part on the discovery of a mutation at codon 181 of HIV reverse transcriptase in non-nucleoside reverse transcriptase inhibitor (nevirapine) treated patient(s) in which the presence of the mutation
15 correlates with a moderate decrease in delavirdine susceptibility and a significant decrease in nevirapine susceptibility and no change in efavirenz susceptibility. The mutation was found in plasma HIV RNA after a period of time following initiation of NNRTI therapy. The
20 development of the codon 181 mutation in HIV RT was found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline. This invention is based in part on
25 the discovery of a mutation at codon 188 of HIV reverse transcriptase in non-nucleoside reverse transcriptase inhibitor (efavirenz) treated patient(s) in which the presence of the mutation correlates with a slight decrease in delavirdine susceptibility and a substantial decrease in
30 nevirapine susceptibility. The mutation was found in plasma HIV RNA after a period of time following initiation of NNRTI therapy. The development of the codon 188 mutation in HIV RT was found to be an indicator of the development of alterations in phenotypic
35 susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline. This invention is based in part on the discovery of a mutation at codon 188 of HIV reverse transcriptase in

patient(s) with no previously reported exposure to non-nucleoside reverse transcriptase inhibitors in which the presence of the mutations correlate with a moderate decrease in delavirdine susceptibility and a substantial
5 decrease in nevirapine susceptibility and a moderate decrease in efavirenz susceptibility. The mutation was found in plasma HIV RNA after a period of time following initiation of anti-retroviral therapy. The development of the codon 138 and 188 mutations in HIV RT was found to be
10 an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline. This invention is based in part on the discovery of a mutation at codon 98 of HIV reverse
15 transcriptase in patient(s) with no previously reported exposure to non-nucleoside reverse transcriptase inhibitors in which the presence of the mutation correlates with slight decrease in delavirdine, nevirapine and efavirenz susceptibility. The mutation was found in plasma HIV RNA
20 after a period of time following initiation of anti-retroviral therapy. The development of the codon 98 mutation in HIV RT was found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with
25 virologic failure and subsequent immunological decline.

This invention is based in part on the discovery of a mutation at codon 98 either alone or in combination with a mutation at codon 190 of HIV reverse transcriptase in
30 patient(s) whose anti-retroviral treatment was unknown in which the presence of the mutations correlate with an increase in delavirdine susceptibility and substantial decrease in both nevirapine and efavirenz susceptibility. The mutations were found in plasma HIV RNA. The
35 development of the mutant at codon 98 in addition to the mutation at codon 190 in HIV RT was found to be an indicator of the development of resistance and ultimately of immunological decline. This invention is based in part

on the discovery of a mutation at codon 181 either alone or in combination with a mutation at codon 98 of HIV reverse transcriptase in non-nucleoside reverse transcriptase inhibitor (delavirdine) treated patient(s) in which the presence of the mutations correlate with an significant decreases in delavirdine susceptibility and a substantial decrease in efavirenz susceptibility. The mutations were found in plasma HIV RNA after a period of time following initiation of therapy. The development of the mutant at codon 98 in addition to the mutation at codon 181 in HIV RT was found to be an indicator of the development of resistance and ultimately of immunological decline. This invention is based in part on the discovery of a mutation at codon 101 either alone or in combination with a mutation at codon 190, for example 190s of HIV reverse transcriptase in non-nucleoside reverse transcriptase inhibitor (efavirenz) treated patient(s) in which the presence of the mutations correlate with no change in delavirdine susceptibility and a substantial decrease in both nevirapine and efavirenz susceptibility. The mutations were found in plasma HIV RNA after a period of time following initiation of therapy. The development of the mutant at codon 101 in addition to the mutation at codon 190, for example 190s in HIV RT was found to be an indicator of the development of resistance and ultimately of immunological decline. This invention is based in part on the discovery of a mutation at codon 108 of HIV reverse transcriptase in patient(s) with no previously reported exposure to non-nucleoside reverse transcriptase inhibitor in which the presence of the mutation correlates with no change in delavirdine susceptibility and a slight decrease in nevirapine susceptibility and no change in efavirenz susceptibility. The mutation was found in plasma HIV RNA after a period of time following initiation of anti-retroviral therapy. The development of the codon 108 mutation in HIV RT was found to be an indicator of the development of alterations in phenotypic

susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline.

This invention is based in part on the discovery of a
5 mutation at codon 101 either alone or in combination with
a mutation at codon 103 and/or 190 of HIV reverse
transcriptase in patients with no previously reported
exposure to non-nucleoside reverse transcriptase inhibitors
in which the presence of the mutations correlate with
10 changes in delavirdine, nevirapine and efavirenz
susceptibility. Specifically, the presence of mutations at
101 and 190, for example 190A, correlates with no change in
delavirdine susceptibility and a substantial decrease in
nevirapine susceptibility and a significant decrease in
15 efavirenz susceptibility. The presence of mutations at 103
and 190 correlates with a moderate decrease in delavirdine
susceptibility, a substantial decrease in nevirapine
susceptibility and a significant decrease in efavirenz
susceptibility. The mutations were found in plasma HIV RNA
20 after a period of time following initiation of anti-
retroviral therapy. The development of the codon 101 and
103 and/or 190 mutations in HIV RT was found to be an
indicator of the development of alterations in phenotypic
susceptibility/resistance which has been associated with
25 virologic failure and subsequent immunological decline.
This invention is based in part on the discovery of a
mutation at codon 106 either alone or in combination with
a mutation at codon 189 and/or 181 and 227 of HIV reverse
transcriptase in non-nucleoside reverse transcriptase
30 inhibitor (nevirapine) treated patient(s) in which the
presence of the mutations correlate with changes in
delavirdine, nevirapine and efavirenz susceptibility.
Specifically, the presence of mutations at 106 and 181
correlates with a significant decrease in delavirdine
35 susceptibility, a substantial decrease in nevirapine
susceptibility and a slight decrease in efavirenz
susceptibility. The presence of mutations at 106 and 189
correlates with a slight decrease in delavirdine

susceptibility, a moderate decrease in nevirapine susceptibility and no change in efavirenz susceptibility. The presence of mutations at 106 and 227 correlates with a slight decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. The presence of mutations at 181 and 227 correlates with an increase in delavirdine susceptibility, a significant decrease in nevirapine susceptibility and an increase in efavirenz susceptibility. The presence of mutations at 106 and 181 and 227 correlates with a moderate decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. The mutations were found in plasma HIV RNA after a period of time following initiation of NNRTI therapy. The development of the codon 106 and 189 and/or 181 and 227 mutations in HIV RT was found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline. This invention is based in part on the discovery of a mutation at codon 103 either alone or in combination with a mutation at codon 100 and/or 188 of HIV reverse transcriptase in non-nucleoside reverse transcriptase inhibitor (nevirapine) treated patient(s) in which the presence of the mutations correlate with changes in delavirdine, nevirapine and efavirenz susceptibility. Specifically, the presence of mutations at 103 and 188 correlates with a substantial decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. The presence of mutations at 100 and 103 correlates with a substantial decrease in delavirdine susceptibility, a moderate decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. The presence of mutations at 103 and 100 and 188 correlates with a substantial decrease in delavirdine susceptibility, a substantial decrease in

- nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. The mutations were found in plasma HIV RNA after a period of time following initiation of NNRTI therapy. The developemnt of the codon 103 and 100
5 and/or 188 mutations in HIV RT was found to be an indicator of the development of alterations in phenotypic susceptibility/resistance which has been associated with virologic failure and subsequent immunological decline.
- 10 In a further embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 225 in combination with mutations at other codons including 103 of HIV RT which correlate with a specific pattern of resistance to antiretroviral
15 therapies and subsequent immunologic decline. In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 236 either alone or in combination with mutations at other codons including 103
20 and/or 181 of HIV RT which correlate with resistance to antiretroviral therapy and immunologic decline. In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 190 (G190S) either alone or in
25 combination with mutation at codon 101 (K101E) of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

- In still another embodiment of the invention, PCR based
30 assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 190 (G190A) either alone or in combination with mutation at codon 103 (K103N) of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

- 35 In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 230 either alone or in

combination with mutation at codon 181 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

- 5 In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect a mutation at codon 181 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

10

In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect a mutation at codon 188 of HIV RT which correlates with resistance to antiretroviral therapy and

- 15 immunologic decline.

- In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 138 either alone or in
20 combination with mutation at codon 188 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

- In yet another embodiment of the invention, PCR based
25 assays, including phenotypic and genotypic assays, may be used to detect a mutation at codon 98 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

- 30 In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 98 either alone or in combination with mutation at codon 190 of HIV RT which correlates with resistance to antiretroviral therapy and
35 immunologic decline.

In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be

used to detect mutations at codon 181 either alone or in combination with mutation at codon 98 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

5

In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 101 either alone or in combination with mutation at codon 190, for example 190s of
10 HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be
15 used to detect a mutation at codon 108 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

In yet another embodiment of the invention, PCR based
20 assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 101 either alone or in combination with mutations at codon 103 and/or 190 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

25

In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be used to detect mutations at codon 106 either alone or in combination with mutations at codon 189 and/or 181 and 227
30 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline.

In yet another embodiment of the invention, PCR based assays, including phenotypic and genotypic assays, may be
35 used to detect mutations at codon 188 either alone or in combination with mutation at codon 100 and /or 103 of HIV RT which correlates with resistance to antiretroviral therapy and immunologic decline. Once mutations at codon

225 and 103 have been detected in a patient undergoing NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 236 and/or 103 and/or 181 have been
5 detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 190 and/or 103 and/or 101 have been detected in a
10 patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 230 and/or 181 have been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic
15 regimen must be considered. Similarly, once a mutation at codon 181 has been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once a
20 mutation at codon 188 has been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 138 and/or 188 have been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic
25 regimen must be considered. Similarly, once a mutation at codon 98 has been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once
30 mutations at codon 98 and/or 190 have been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 181 and/or 98 have been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic
35 regimen must be considered. Similarly, once mutations at codon 101 and/or 190, for example 190S, have been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once a mutation at codon 108 has

been detected in a patient underfoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 101 and/or 103 and/or 190, for example 190A, have
5 been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 106 and/or 189 and/or 181 and/or 227 have been detected in a patient undergoing certain NNRTI
10 antiretroviral therapy, an alteration in the therapeutic regimen must be considered. Similarly, once mutations at codon 188 and/or 100 and/or 103 have been detected in a patient undergoing certain NNRTI antiretroviral therapy, an alteration in the therapeutic regimen must be considered.
15 The timing at which a modification of the therapeutic regimen should be made, following the assessment of the antiretroviral therapy using PCR based assays, may depend on several factors including the patient's viral load, CD4 count, and prior treatment history.

20

In another aspect of the invention there is provided a method for assessing the effectiveness of a non-nucleoside reverse transcriptase antiretroviral drug comprising: (a) introducing a resistance test vector comprising a patient-
25 derived segment and an indicator gene into a host cell; (b) culturing the host cell from step (a); (c) measuring expression of the indicator gene in a target host cell wherein expression of the indicator gene is dependent upon the patient derived segment; and (d) comparing the
30 expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the NNRTI anti-HIV drug, wherein a test concentration of the NNRTI, anti-HIV drug is presented at steps (a) - (c); at steps (b) - (c);
35 or at step (c).

This invention also provides a method for assessing the effectiveness of non-nucleoside reverse transcriptase

antiretroviral therapy in a patient comprising: (a) developing a standard curve of drug susceptibility for an NNRTI anti-HIV drug; (b) determining NNRTI anti-HIV drug susceptibility in the patient using the susceptibility test
5 described above; and (c) comparing the NNRTI anti-HIV drug susceptibility in step (b) with the standard curve determined in step (a), wherein a decrease in NNRTI anti-HIV susceptibility indicates development of anti-HIV drug resistance in the patient.

10

This invention also provides a method for evaluating the biological effectiveness of a candidate HIV antiretroviral drug compound comprising: (a) introducing a resistance test
15 vector comprising a patient-derived segment and an indicator gene into a host cell; (b) culturing the host cell from step (a); (c) measuring expression of the indicator gene in a target host cell wherein expression of the indicator gene is dependent upon the patient derived
20 gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound, wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) - (c);
25 or at step (c).

The expression of the indicator gene in the resistance test vector in the target cell is ultimately dependent upon the action of the patient-derived segment sequences. The
30 indicator gene may be functional or non-functional.

In another aspect this invention is directed to antiretroviral drug susceptibility and resistance tests for HIV/AIDS. Particular resistance test vectors of the
35 invention for use in the HIV/AIDS antiretroviral drug susceptibility and resistance test are identified.

In yet another aspect this invention provides for the identification and assessment of the biological effectiveness of potential therapeutic antiretroviral compounds for the treatment of HIV and/or AIDS. In another
5 aspect, the invention is directed to a novel resistance test vector comprising a patient-derived segment further comprising one or more mutations on the RT gene and an indicator gene.

Brief Description of the Drawings

Fig. 1

Resistance Test Vector. A diagrammatic representation of the resistance test vector comprising a patient derived
5 segment and an indicator gene.

Fig. 2

Two Cell Assay. Schematic Representation of the Assay. A resistance test vector is generated by cloning the patient-
10 derived segment into an indicator gene viral vector. The resistance test vector is then co-transfected with an expression vector that produces amphotropic murine leukemia virus (MLV) envelope protein or other viral or cellular proteins which enable infection. Pseudotyped viral
15 particles are produced containing the protease (PR) and the reverse transcriptase (RT) gene products encoded by the patient-derived sequences. The particles are then harvested and used to infect fresh cells. Using defective PR and RT sequences it was shown that luciferase activity
20 is dependent on functional PR and RT. PR inhibitors are added to the cells following transfection and are thus present during particle maturation. RT inhibitors, on the other hand, are added to the cells at the time of or prior to viral particle infection. The assay is performed in the
25 absence of drug and in the presence of drug over a wide range of concentrations. The amount of luciferase is determined and the percentage (%) inhibition is calculated at the different drug concentrations tested.

30 Fig. 3

Examples of phenotypic drug susceptibility profiles. Data are analyzed by plotting the percent inhibition of luciferase activity vs. \log_{10} concentration (μM). This plot is used to calculate the drug concentration that is
35 required to inhibit virus replication by 50% (IC_{50}) or by 95% (IC_{95}). Shifts in the inhibition curves towards higher drug concentrations are interpreted as evidence of drug resistance. Three typical curves for a nucleoside reverse

transcriptase inhibitor (AZT), a non-nucleoside reverse transcriptase inhibitor (delavirdine), and a protease inhibitor (ritonavir) are shown. A reduction in drug susceptibility (resistance) is reflected in a shift in the
5 drug susceptibility curve toward higher drug concentrations (to the right) as compared to a baseline (pre-treatment) sample or a drug susceptible virus control, such as PNL4-3 or HXB-2, when a baseline sample is not available.

10 Fig. 4

Phenotypic drug susceptibility and resistance profile: patient 487. A PCR-based phenotypic susceptibility assay was carried out giving the phenotypic drug susceptibility and resistance profile showing increased resistance to both
15 delavirdine and nevirapine. This is an example of the first pattern of NNRTI susceptibility/resistance. Evaluation of this virus from plasma showed HIV reverse transcriptase having mutations at codons 184 (M184V) associated with 3TC resistance and at 103 (K103N)
20 associated with both delavirdine and nevirapine resistance.

Fig. 5

Phenotypic drug susceptibility and resistance profile of site directed reverse transcriptase mutants. A PCR-based
25 phenotypic susceptibility assay was carried out giving the phenotypic drug susceptibility and resistance profile for site directed mutants having mutations at codons 103 and 181 (K103N; Y181C) demonstrating resistance to both delavirdine and nevirapine. The double mutant demonstrates
30 the additive effect of both mutations resulting in a further increase in resistance.

Fig. 6

Phenotypic drug susceptibility and resistance profile:
35 Patient 268. A PCR-based phenotypic susceptibility assay was carried out giving the phenotypic drug susceptibility and resistance profile showing the evaluation of virus from plasma with HIV reverse transcriptase having phenotypic

resistance to delavirdine but not nevirapine. This is an example of the second pattern of NNRTI susceptibility/resistance. This patient virus is resistant to all of the protease inhibitors tested and also has significant resistance to AZT and 3TC and shows slight shifts in susceptibility to ddC, ddI, and d4T. Evaluation of this virus from plasma using a PCR and sequencing based genotypic assay showed HIV reverse transcriptase having mutations at codons 103 and 236 (K103N; P236L). The P236L mutation was previously reported to cause delavirdine resistance and nevirapine hypersensitivity (Dueweke TJ et al. (1993) *Proc Natl Acad Sci* 90, 4713-4717). However, in this patient sample, while there was delavirdine resistance nevirapine susceptibility was the same as wild type.

15

Fig. 7

Phenotypic drug susceptibility and resistance profile of site-directed reverse transcriptase mutant (P236L). A PCR-based phenotypic susceptibility assay was carried out giving the phenotypic drug susceptibility and resistance profile showing the susceptibility to delavirdine and nevirapine of the P236L site-directed mutagenesis mutant. This result is identical to that observed in the patient virus sample shown in Figure 6. The next two panels show the K103N site-directed mutagenesis mutant and the two panels below show the double mutant K103N + P236L. The P236L mutation is additive to the K103N causing severe resistance to delavirdine while having no effect on nevirapine resistance due to K103N. The right side of the figure shows a similar result when the P236L mutation is added to the Y181->C mutation.

Fig. 8A

Phenotypic Drug Susceptibility and Resistance Profile: Patients 302. This is one example of the third pattern of NNRTI susceptibility/resistance. Phenotypic analysis of the patient virus demonstrated reduced susceptibility to both delavirdine and nevirapine. This pattern is

characterized by a larger reduction of nevirapine susceptibility compared to the reduction of delavirdine susceptibility. Genotypic analysis of the patient virus demonstrated the presence of the RT mutations K103N
5 associated with nevirapine and delavirdine resistance and P225H.

Fig. 8B

Phenotypic Drug Susceptibility and Resistance Profile:
10 Patients 780. This is a second example of the third pattern of NNRTI susceptibility/resistance. Phenotypic analysis of the patient virus demonstrated reduced susceptibility to both delavirdine and nevirapine. This pattern is characterized by a larger reduction of
15 nevirapine susceptibility compared to the reduction of delavirdine susceptibility. Genotypic analysis of the patient virus demonstrated the presence of the RT mutations K103N associated with nevirapine and delavirdine resistance and P225H.

20

Fig. 8C

Phenotypic Drug Susceptibility and Resistance Profile:
Individual Virus Clones of Patient 302. Genotypic analysis of individual virus clones from patient 302 revealed
25 viruses containing the K103N mutation without the P225H mutation (K103N, I135M, R211K) and viruses containing the K103N mutation with the P225H mutation (K103N, P225H). Phenotypic characterization of these virus clones indicates that the P225H mutation reduces the amount delavirdine
30 resistance associated with the K103N mutation (compare bottom panels), but does not alter the amount of nevirapine resistance associated with the K103N mutation (compare top panels).

35 Fig. 8D

Phenotypic Drug Susceptibility and Resistance Profile: Site Directed Reverse Transcriptase Mutants. Phenotypic characterization of a virus containing the site directed RT

mutation P225H indicates that this mutation increases susceptibility to delavirdine, but not nevirapine (compare top panels). Phenotypic characterization of a virus containing the site directed RT mutations P225H plus K013N or P225H plus Y181C indicate that the P225H mutation decreases the amount of delavirdine resistance associated with either K103N or Y181C, but does not decrease the amount of nevirapine resistance associated with K103N or Y181C. to delavirdine, but not nevirapine (compare corresponding middle and bottom panels).

Fig. 9A

Phenotypic Drug Susceptibility and Resistance Profile: Patients 644. This is one example of the fourth pattern of NNRTI susceptibility and resistance. Phenotypic analysis of the patient virus demonstrated by a large reduction in susceptibility to nevirapine, but not delavirdine. Genotypic analysis of the patient virus demonstrated the presence of the RT mutations G190S, as well as the K101E mutation associated with reductions in susceptibility to atevirdine, DMP266, L-697,661 and UC-10,38,57 (Schinazi, Mellors, Larder resistance table).

Fig. 9B

Phenotypic Drug Susceptibility and Resistance Profile: Site Directed Reverse Transcriptase Mutants. Phenotypic characterizations of viruses containing either site directed RT mutations G190A, or G190S indicate that these mutations greatly reduce susceptibility to nevirapine, and slightly increase susceptibility to delavirdine (compare top panels).

Detailed Description of the Invention

The present invention relates to methods of monitoring the clinical progression of HIV infection in patients receiving antiretroviral therapy, particularly non-nucleoside reverse transcriptase inhibitor antiretroviral therapy.

In one embodiment, the present invention provides for a method of assessing the effectiveness of antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at one or more positions in the RT. The mutation(s) correlate positively with alterations in phenotypic susceptibility/resistance.

In a specific embodiment, the invention provides for a method of assessing the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon 225 and 103. This invention established, using a phenotypic susceptibility assays, that mutations at codon 225 either alone or in combination with a mutation at codon 103 of HIV reverse transcriptase are correlated with an increase in delavirdine susceptibility, little or no change in nevirapine susceptibility and little or no change in efavirenz susceptibility.

In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 236 and 103 and/or 181. This invention established, using a phenotypic susceptibility assay, that mutations at codon 236 either alone or in combination with

a mutation at codon 103 and/or 181 of HIV reverse transcriptase are correlated with a decrease in delavirdine susceptibility (increased resistance) and no change in nevirapine susceptibility. The 236 mutation alone or on a
5 Y181C background has no effect on efavirenz susceptibility but restores a significant portion of the loss of susceptibility caused by a 103N mutation.

In another specific embodiment, the invention provides for
10 a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at
15 codon(s) 230 and/or 181. This invention established, using a phenotypic susceptibility assay, that mutations at codon 230 either alone or in combination with a mutation at codon 181 of HIV reverse transcriptase are correlated with a significant decrease in delavirdine susceptibility
20 (increased resistance), significant decrease in nevirapine susceptibility.

In another specific embodiment, the invention provides for
a method of evaluating the effectiveness of NNRTI
25 antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon 181. This invention established, using a phenotypic
30 susceptibility assay, that a mutation at codon 181 of HIV reverse transcriptase is correlated with a moderate decrease in delavirdine susceptibility (increased resistance), significant decrease in nevirapine susceptibility and no change in efavirenz susceptibility.

35

In another specific embodiment, the invention provides for
a method of evaluating the effectiveness of NNRTI
antiretroviral therapy of a patient comprising (i)

collecting a biological sample from an HIV-infected patient, and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon 188. This invention established, using a phenotypic susceptibility assay, that a mutation at codon 188 of HIV reverse transcriptase are correlated with a slight decrease in delavirdine susceptibility (increased resistance), a substantial decrease in nevirapine susceptibility and a significant decrease in efavirenz susceptibility.

10

In other specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 138 and/or 188. This invention established, using a phenotypic susceptibility assay, that mutations at codon 138 either alone or in combination with a mutation at codon 188 of HIV reverse transcriptase are correlated with a moderate decrease in delavirdine susceptibility (increased resistance), a substantial decrease in nevirapine susceptibility and a moderate decrease in efavirenz susceptibility.

25

In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 98. This invention established, using a phenotypic susceptibility assays, that mutations at codon 98 of HIV reverse transcriptase are correlated with a slight decrease in delavirdine susceptibility (increase resistance), a slight decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility.

35

In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 98 and/or 190. This invention established, using a phenotypic susceptibility assay, that mutations at codon 98 either alone or in combination with a mutation at codon 190 of HIV reverse transcriptase are correlated with an increase in delavirdine susceptibility (decreased resistance), a substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. In other specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 181 and/or 98. This invention established, using a phenotypic susceptibility assay, that mutations at codon 181 either alone or in combination with a mutation at codon 98 of HIV reverse transcriptase are correlated with a significant decrease in delavirdine susceptibility (increased resistance), a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient ; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 101 and/or 190, for example 190S. This invention established, using a phenotypic susceptibility assay, that mutations at codon 101 either alone or in combination with a mutation at codon 190 of HIV reverse transcriptase are correlated with no change in delavirdine susceptibility (wild-type), a

substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 108. This invention established, using a phenotypic susceptibility assay, that a mutation at codon 108 of HIV reverse transcriptase are correlated with a no change in delavirdine susceptibility (wild-type), a slight decrease in nevirapine susceptibility and no change in efavirenz susceptibility. In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 101 and 103 and/or 190. This invention established, using a phenotypic susceptibility assay, that mutations at codon 101 either alone or in combination with a mutation at codon 103 and/or 190 of HIV reverse transcriptase are correlated with a either no change (101 and 190) or a moderate decrease (103 and 190, for example 190A) in delavirdine susceptibility (increased resistance), a substantial decrease in nevirapine susceptibility and a significant decrease in efavirenz susceptibility.

30

In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 106 and/or 189 and/or 181 and/or 227. This invention established, using a phenotypic susceptibility

assay, that mutations at codon 106 either alone or in combination with a mutation at codon 189 and/or 181 and/or 227 of HIV reverse transcriptase are correlated with changes in delavirdine, nevirapine and efavirenz susceptibility. Specifically, the presence of mutations at 106 and 181 correlates with a significant decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. The presence of mutations at 106 and 189 correlates with a slight decrease in delavirdine susceptibility, a moderate decrease in nevirapine susceptibility and no change in efavirenz susceptibility. The presence of mutations at 106 and 227 correlates with a slight decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. The presence of mutations at 181 and 227 correlates with an increase in delavirdine susceptibility, a significant decrease in nevirapine susceptibility and an increase in efavirenz susceptibility. The presence of mutations at 106 and 181 and 227 correlates with a moderate decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. In another specific embodiment, the invention provides for a method of evaluating the effectiveness of NNRTI antiretroviral therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; and (ii) determining whether the biological sample comprises nucleic acid encoding HIV RT having a mutation at codon(s) 188 and 100 and/or 103. This invention established, using a phenotypic susceptibility assay, that mutations at codon 188 either alone or in combination with a mutation at codon 100 and/or 103 of HIV reverse transcriptase are correlated changes in delavirdine, nevirapine and efavirenz susceptibility. Specifically, the presence of mutations at 103 and 188 correlates with a substantial decrease in delavirdine susceptibility, a substantial decrease in nevirapine

susceptibility and a substantial decrease in efavirenz susceptibility. The presence of mutations at 100 and 103 correlates with a substantial decrease in delavirdine susceptibility, a moderate decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. The presence of mutations at 103 and 100 and 188 correlates with a substantial decrease in delavirdine susceptibility, a substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. Under the foregoing circumstances, the phenotypic susceptibility/resistance profile and genotypic profile of the HIV virus infecting the patient has been altered reflecting some change in the response to the antiretroviral agent. In the case on NNRTI antiretroviral therapy, the HIV virus infecting the patient may be resistant to one or more but not another of the NNRTIs as described herein. It therefore may be desirable after detecting the mutation, to either increase the dosage of the antiretroviral agent, change to another antiretroviral agent, or add one or more additional antiretroviral agents to the patient's therapeutic regimen. For example, if the patient was being treated with efavirenz (DMP-266) when the 225 mutation arose, the patient's therapeutic regimen may desirably be altered by either (i) changing to a different NNRTI antiretroviral agent, such as delavirdine or nevirapine and stopping efavirenz treat; or (ii) increasing the dosage of efavirenz; or (iii) adding another antiretroviral agent to the patient's therapeutic regimen. The effectiveness of the modification in therapy may be evaluated by monitoring viral burden such as by HIV RNA copy number. A decrease in HIV RNA copy number correlates positively with the effectiveness of a treatment regiment.

The phrase "correlates positively," as used herein, indicates that a particular result renders a particular conclusion more likely than other conclusions.

- 5 Another preferred, non-limiting, specific embodiment of the invention is as follows: A method of assessing the effectiveness of NNRTI therapy of a patient comprising (i) collecting a biological sample from an HIV-infected patient; (ii) amplifying the HIV-encoding RNA in the biological
10 sample by converting the RNA to cDNA and amplifying HIV sequences using HIV primers that result in a PCR product that comprises that RT gene; (iii) performing PCR using primers that result in PCR products comprising wild type or mutant 225 and 103 codons; and (iv) determining, via the
15 products of PCR, the presence or absence of a mutation at codon 225 or 103 or both. Yet another preferred, non-limiting specific embodiment, of the invention is as follows: A method of assessing the effectiveness of NNRTI therapy of a patient comprising (I) collecting a plasma
20 sample from an HIV-infected patient; (ii) amplifying the HIV-encoding RNA in the plasma sample by converting the RNA to cDNA and amplifying HIV sequences using HIV primers that result in a PCR product that comprises the RT gene; (iii) performing PCR using primers that result in PCR products
25 comprising the wild type or mutations at codons 103 and/or 181 and 236; and (iv) determining, via the products of PCR, the presence or absence of a mutation at codon 236 and 103 and/or 181.
- 30 Yet another preferred, non-limiting specific embodiment, of the invention is as follows: A method of assessing the effectiveness of NNRTI therapy of a patient comprising (I) collecting a plasma sample from an HIV-infected patient; (ii) amplifying the HIV-encoding RNA in the plasma sample by
35 converting the RNA to cDNA and amplifying HIV sequences

using HIV primers that result in a PCR product that comprises that RT gene: (iii) performing PCR using primers that result in PCR products comprising the wild type or mutations at codon 101 and 190 (G190S); and (iv) 5 determining, via the products of PCR, the presence or absence of a mutation at codon 190 (G190S) and 101.

Yet another preferred, non-limiting specific embodiment, of the invention is as follows: A method of assessing the 10 effectiveness of NNRTI therapy of a patient comprising (i) collecting a plasma sample from an HIV-infected patient; (ii) amplifying the HIV-encoding RNA in the plasma sample by converting the RNA to cDNA and amplifying HIV sequences using HIV primers that result in a PCR product that 15 comprises the RT gene; (iii) performing PCR using primers that result in PCR products comprising the wild type or mutations at codon 103 and 190 (G190A) and (iv) determining, via the products of PCR, the presence or absence of a mutation at codon 190 (G190A) and 103. Yet another 20 preferred, non-limiting specific embodiment, of the invention is as follows: A method of assessing the effectiveness of NNRTI therapy of a patient comprising (i) collecting a plasma sample from an HIV-infected patient; (ii) amplifying the HIV-encoding RNA in the plasma sample by 25 converting the RNA to cDNA and amplifying HIV sequences using HIV primers that result in a PCR product that comprises the RT gene; (iii) performing PCR using primers that result in PCR products comprising the wild type or mutations at codon 230 and 181, and (iv) determining, via 30 the products of PCR, the presence or absence of a mutation at codon 230 and 181.

Yet another preferred, non-limiting specific embodiment, of the invention is as follows: A method of assessing the 35 effectiveness of NNRTI therapy of a patient comprising (i)

collecting a plasma sample from an HIV-infected patient;
(ii) amplifying the HIV-encoding RNA in the plasma sample by
converting the RNA to cDNA and amplifying HIV sequences
using HIV primers that result in a PCR product that
5 comprises the RT gene; (iii) performing PCR using primers
that result in PCR products comprising the wild type or
mutation at 181; and (iv) determining, via the products of
PCR, the presence or absence of a mutation at codon 181.
Yet another preferred, non-limiting specific embodiment, of
10 the invention is as follows: A method of assessing the
effectiveness of NNRTI therapy of a patient comprising (i)
collecting a plasma sample from an HIV-infected patient;
(ii) amplifying the HIV-encoding RNA in the plasma sample by
converting the RNA to cDNA and amplifying HIV sequences
15 using HIV primers that result in a PCR product that
comprises the RT gene; (iii) performing PCR using primers
that result in PCR products comprising the wild type or
mutation at codon 188; and (iv) determining, via the
products of PCR, the presence or absence of a mutation at
20 codon 188. Yet another preferred, non-limiting specific
embodiment, of the invention is as follows: A method of
assessing the effectiveness of NNRTI therapy of a patient
comprising (i) collecting a plasma sample from an HIV-
infected patient; (ii) amplifying the HIV-encoding RNA in
25 the plasma sample by converting the RNA to cDNA and
amplifying HIV sequences using HIV primers that result in a
PCR product that comprises the RT gene; (iii) performing PCR
using primers that result in PCR products comprising the
wild type or mutations at codon 138 and 188; and (iv)
30 determining, via the products of PCR, the presence or
absence of a mutation at codon 138 and 188.

Yet another preferred, non-limiting specific embodiment, of
the invention is as follows: A method of assessing the
35 effectiveness of NNRTI therapy of a patient comprising (i)

collecting a plasma sample from an HIV-infected patient;
(ii) amplifying the HIV-encoding RNA in the plasma sample by
converting the RNA to cDNA and amplifying HIV sequences
using HIV primers that result in a PCR product that
5 comprises the RT gene; (iii) performing PR using primers
that result in PCR products comprising the wild type or
mutation at codon 98 and (iv) determining, via the products
of PCR, the presence or absence of a mutation at codon 98.
Yet another preferred, non-limiting specific embodiment, of
10 the invention is as follows: A method of assessing the
effectiveness of NNRTI therapy of a patient comprising (i)
collecting a plasma sample from an HIV-infected patient;
(ii) amplifying the HIV-encoding RNA in the plasma sample by
converting the RNA to cDNA and amplifying HIV sequences
15 using HIV primers that result in a PCR product that
comprises the RT gene; (iii) performing PCR using primers
that result in PCR products comprising the wild type or
mutations at codon 98 and 190; and (iv) determining, via the
products of PCR, the presence or absence of a mutation at
20 codon 190 and 98. Yet another preferred, non-limiting
specific embodiment, of the invention is as follows: A
method of assessing the effectiveness of NNRTI therapy of a
patient comprising (i) collecting a plasma sample from an
HIV-infected patient; (ii) amplifying the HIV-encoding RNA
25 in the plasma sample by converting the RNA to cDNA and
amplifying HIV sequences using HIV primers that result in a
PCR product that comprises the RT gene; (iii) performing PCR
using primers that result in PCR products comprising the
wild type or mutations at codon 98 and 181; and (iv)
30 determining, via the products of PCR, the presence or
absence of a mutation at codon 98 and 181.

Yet another preferred, non-limiting specific embodiment, of
the invention is as follows: A method of assessing the
35 effectiveness of NNRTI therapy of a patient comprising (i)

collecting a plasma sample from an HIV-infected patient;
(ii) amplifying the HIV-encoding RNA in the plasma sample by
converting the RNA to cDNA and amplifying HIV sequences
using HIV primers that result in a PCR product that
5 comprises the RT gene; (iii) performing PCR using primers
that result in PCR products comprising the wild type or
mutations at codon 101 and 190; and (iv) determining, via
the products of PCR, the presence or absence of a mutation
at codon 190, for example 190S and 101.

10

Yet another preferred, non-limiting specific embodiment, of
the invention is as follows: A method of assessing the
effectiveness of NNRTI therapy of a patient comprising (i)
collecting a plasma sample from an HIV-infected patient;
15 (ii) amplifying the HIV-encoding RNA in the plasma sample by
converting the RNA to cDNA and amplifying HIV sequences
using HIV primers that result in a PCR product that
comprises the RT gene; (iii) performing PCR using primers
that result in PCR products comprising the wild type or a
20 mutation at codon 108; and (iv) determining, via the
products of PCR, the presence or absence of a mutation at
codon 108. Yet another preferred, non-limiting specific
embodiment, of the invention is as follows: A method of
assessing the effectiveness of NNRTI therapy of a patient
25 comprising (i) collecting a plasma sample from an HIV-
infected patient; (ii) amplifying the HIV-encoding RNA in
the plasma sample by converting the RNA to cDNA and
amplifying HIV sequences using HIV primers that result in a
PCR product that comprises the RT gene; (iii) performing PCR
30 using primers that result in PCR products comprising the
wild type or a mutation at codon 101 and 103 and 190 and
(iv) determining, via the products of PCR, the presence or
absence of a mutation at codon 101 and 103 and 190, for
example 190A.

35

Yet another preferred, non-limiting specific embodiment, of the invention is as follows: A method of assessing the effectiveness of NNRTI therapy of a patient comprising (i) collecting a plasma sample from an HIV-infected patient; 5 (ii) amplifying the HIV-encoding RNA in the plasma sample by converting the RNA to cDNA and amplifying HIV sequences using HIV primers that result in a PCR product that comprises that RT gene; (iii) performing PCR using primers that result in PCR products comprising the wild type or 10 mutations at codon 106 and and 189 and 181 and 227 and (iv) determining, via the products of PCR, the presence or absence of a mutation at codon 106 and 189 and 181 and 227.

Yet another preferred, non-limiting specific embodiment, of 15 the invention is as follows: A method of assessing the effectiveness of NNRTI therapy of a patient comprising (i) collecting a plasma sample from an HIV-infected patient; (ii) amplifying the HIV-encoding RNA in the plasma sample by converting the RNA to cDNA and amplifying HIV sequences 20 using HIV primers that result in a PCR product that comprises the RT gene; (iii) performing PCR using primers that result in PCR products comprising the wild type or mutations at codon 188 and 100 and 103 and (iv) determining, via the products of PCR, the presence or absence of a 25 mutation at codon 188 and 100 and 103. The presence of the mutation at codon 225 and 103 of HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy may require alteration, since as shown by this invention mutation at codon 103 reduces susceptibility which 30 susceptibility can in part be restored by mutation at codon 225. Using the methods of this invention change in the NNRTI therapy would be indicated. Similarly, using the means and methods of this invention the presence of the mutation at codon 236 and 103 and/or 181 of the HIV RT 35 indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly,

using the means and methods of this invention the presence of the mutation at codon 190 (G190A) and 103 (K103N) of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly,

5 using the means and methods of this invention the presence of the mutation at codon 190 (G190S) and 101 (K101E) of the HIV RT indicate that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly,

10 using the means and methods of this invention the presence of the mutation at codon 230 and 181 of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence of the a mutation at codon 181 of the HIV RT indicates that the effectiveness of

15 the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence of the mutation at codon 188 of the HIV RT indicates that the effectiveness of the current of prospective NNRTI therapy has been diminished. Similarly,

20 using the means and methods of this invention the presence of the mutation at codon 138 and 188 of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence of the mutation at

25 codon 98 of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence of the mutation at codon 98 and 190 of the HIV RT indicates that the effectiveness of the

30 current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence of the mutation at codon 181 and 98 of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly,

35 using the means and methods of this invention the presence of the mutation at codon 101 and 190, for example 190S, of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished.

Similarly, using the means and methods of this invention the presence of a mutation at codon 108 of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence of the mutation at 5 101 and 103 and 190, for example 190A, of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of this invention the presence 10 of the mutation at codon 106 and 189 and 181 and 227 of the HIV RT indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished. Similarly, using the means and methods of the invention the presence of the mutation at codon 188 and 100 and 103 of the HIV RT 15 indicates that the effectiveness of the current or prospective NNRTI therapy has been diminished.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of evaluating the 20 effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 236 and 103 25 and/or 181. Using the phenotypic susceptibility assay, it was observed that the presence of the three mutations correlates positively with delavirdine resistance. Using the phenotypic susceptibility assay, it was observed that the presence of the three mutations correlates positively 30 with nevirapine resistance. In another embodiment, the mutated codon 236 of HIV RT encodes leucine (L). In a further embodiment, the reverse transcriptase has a mutation at codon 103, a mutation at codon 181 or a combination thereof in addition to the mutation at codon 236 of HIV RT. 35 In a still further embodiment, the mutated codon 103 encodes an asparagine (N) and the mutated codon at 181 encodes a cysteine (C).

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from
5 an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 225 and 103. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 225
10 alone or in combination with a mutation at codon 103 of HIV RT cause an increase in delavirdine susceptibility while having no effect on nevirapine susceptibility. In yet another embodiment, the mutated codon 225 codes for a histidine, codon 230 codes for a leucine and codon 181 codes
15 for a cysteine.

This invention provides a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from
20 an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 181. Using the phenotypic susceptibility assay it was observed that the presence of mutations at codon 181 correlates positively
25 with a moderate decrease in delavirdine susceptibility and a significant decrease in nevirapine susceptibility and no change in efavirenz susceptibility. In an embodiment, the mutated codon 181 for a isoleucine.

30 This invention provides a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV
35 reverse transcriptase having a mutation at codon 188. Using the phenotypic susceptibility assay it was observed that

the presence of mutations at codon 188 correlates positively with a slight decrease in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and significant decrease in efavirenz susceptibility. In an embodiment, the mutated codon 188 codes for a cysteine, histidine, or leucine.

This invention provides a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 190. Using the phenotypic susceptibility assay it was observed that the presence of mutations at codon 190 correlates positively with a slight increase in delavirdine susceptibility and a large decrease in nevirapine susceptibility. In an embodiment, the mutated codon 190 codes for an alanine or a serine.

20

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 230 and 181. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 230 alone or in combination with a mutation at codon 181 of HIV RT causes a significant decrease in delavirdine susceptibility and a significant decrease in nevirapine susceptibility.

30

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from
5 an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 138 and 188. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 138
10 alone or in combination with a mutation at codon 188 of HIV RT causes a moderate decrease in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and a moderate decrease in efavirenz susceptibility. In yet another embodiment, the mutated codon 138 codes for a
15 alanine and codon 188 codes for a leucine.

This invention provides a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from
20 an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 98. Using the phenotypic susceptibility assay it was observed that the presence of mutations at codon 98 correlates positively with
25 a slight decrease in delavirdine susceptibility and a slight decrease in delavirdine susceptibility and a slight decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. In an embodiment, the mutated codon 98 codes for glycine.

30

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from
35 an HIV-infected patient; and (b) determining whether the

biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 98 and 190. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 98 alone or in
5 combination with a mutation at codon 190 of HIV RT causes an increase in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. In yet another
10 embodiment, the mutated codon 190 codes for a serine and codon 98 for a glycine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of HIV-infected
15 patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 181 and 98. Using the phenotypic susceptibility assay, it was observed
20 that the presence of the mutations at codons 181 alone or in combination with a mutation at codon 98 of HIV RT causes a significant decrease in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. In yet another
25 embodiment, the mutated codon 98 codes for a glycine and codon 181 codes for a cysteine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the
30 effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 101 and
35 190, for example 190S. Using the phenotypic susceptibility

assay, it was observed that the presence of the mutations at codons 101 alone or in combination with a mutation at codon 190 of HIV RT causes no change in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and
5 a substantial decrease in efavirenz susceptibility. In yet another embodiment, the mutated codon 190 codes for a serine and codon 101 codes for a glutamine acid.

This invention provides a method of assessing the
10 effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 108. Using
15 the phenotypic susceptibility assay it was observed that the presence of mutations at codon 108 correlates positively with no change in delavirdine susceptibility and a slight decrease in nevirapine susceptibility and no change in efavirenz susceptibility. In an embodiment, the mutated
20 codon 108 codes for a isoleucine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected
25 patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 101 and 190, for example 190A. Using the phenotypic susceptibility
30 assay, it was observed that the presence of the mutations at codons 101 alone or in combination with a mutation at codon 190 of HIV RT causes no change in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and a significant decrease in efavirenz susceptibility. In yet

another embodiment, the mutated codon 190 codes for a glycine and codon 101 codes for a glutamine acid.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 103 and 190. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 103 alone or in combination with a mutation at codon 190 of HIV RT causes a moderate decrease in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and a significant decrease in efavirenz susceptibility. In yet another embodiment, the mutated codon 190 codes for a alanine and codon 103 codes for a asparagine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 106 and 181. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 106 alone or in combination with a mutation at codon 181 of HIV RT causes a significant decrease in delvaridine susceptibility and a substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. In yet another embodiment, the mutated codon 106 codes for a alanine and codon 181 codes for a cysteine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from
5 an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 106 and 189. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 106
10 alone or in combination with a mutation at codon 189 of HIV RT causes a slight decrease in delavirdine susceptibility and a moderate decrease in nevirapine susceptibility and no change in efavirenz susceptibility. In yet another embodiment, the mutated codon 189 codes for a leucine and a
15 codon 106 codes for a alanine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected
20 patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 106 and 227. Using the phenotypic susceptibility assay, it was
25 observed that the presence of the mutations at codons 106 alone or in combination with a mutation at codon 227 of HIV RT causes a slight decrease in delavirdine susceptibility and a substantial decrease in nevirapine susceptibility and a slight decrease in efavirenz susceptibility. In yet
30 another embodiment, the mutated codon 227 codes for a leucine and codon 106 codes for a alanine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the
35 effectiveness of antiretroviral therapy of an HIV-infected

patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 181 and 227. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 181 alone or in combination with a mutation at codon 227 of HIV-RT causes an increase in delavirdine susceptibility and a significant decrease in nevirapine susceptibility and an increase in efavirenz susceptibility.

In yet another embodiment, the mutated codon 227 codes for a leucine and codon 181 codes for cysteine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 106 and 181 and 227. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 106 alone or in combination with a mutation at codon 181 and 227 of HIV RT causes a moderate decrease in delavirdine susceptibility and a slight decrease in efavirenz susceptibility.

In yet another embodiment, the mutated codon 106 codes for an alanine, codon 181 codes for a cysteine and codon 227 codes for a leucine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected

patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 103 and
5 188. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 103 alone or in combination with a mutation at codon 188 of HIV RT causes a substantial decrease in delavirdine
10 susceptibility and a substantial decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility. In yet another embodiment, the mutated codon 188 codes for a leucine and codon 103 codes for a asparagine.

15 Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the
20 biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 100 and 103. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 100 alone or in combination with a mutation at codon 103 of HIV
25 RT causes a substantial decrease in delavirdine susceptibility and a moderate decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility.

30 In yet another embodiment, the mutated codon 100 codes for a isoleucine, codon 103 codes for a asparagine.

Another preferred, non-limiting, specific embodiment of the invention is as follows: a method of assessing the
35 effectiveness of antiretroviral therapy of an HIV-infected

patient comprising: (a) collecting a biological sample from an HIV-infected patient; and (b) determining whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 100 and 103 and 188. Using the phenotypic susceptibility assay, it was observed that the presence of the mutations at codons 100 alone or in combination with a mutation at codon 103 and 188 of HIV RT causes a substantial decrease in delavirdine susceptibility and a moderate decrease in nevirapine susceptibility and a substantial decrease in efavirenz susceptibility.

In yet another embodiment, the mutated codon 100 codes for a isoleucine, codon 103 codes for a asparagine and codon 188 codes for a leucine.

This invention also provides the means and methods to use the resistance test vector comprising an HIV gene further comprising an NNRTI mutation for drug screening. More particularly, the invention describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 225 and 103 for drug screening. The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 236 and 103 and/or 181. The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 190 (G190A) and 103 (K103N). The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 190 (G190S) and 101 (K101E).

The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 230 and 181.

The invention also describes the resistance test vector comprising the HIV reverse transcriptase having a mutation at codon 181.

5 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having a mutation at codon 188.

10 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 138 and 188.

15 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having a mutation at 98.

The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 98 and 190.

20 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 181 and 98.

25 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 101 and 190, for example 190S.

30 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having a mutation at codon 108.

35 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 101 and 103 and/or 190, for example 190A.

The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 106 and 189 and/or 181 and/or 227.

- 5 The invention also describes the resistance test vector comprising the HIV reverse transcriptase having mutations at codons 188 and 100 and/or 103.

- 10 The invention further relates to novel vectors, host cells and compositions for isolation and identification of the non-nucleoside HIV-1 reverse transcriptase inhibitor resistance mutant and using such vectors, host cells and compositions to carry out anti-viral drug screening. This invention also relates to the screening of candidate drugs
15 for their capacity to inhibit said mutant.

EXAMPLE 1: Phenotypic Drug Susceptibility and Resistance Test Using Resistance Test Vectors

Phenotypic drug susceptibility and resistance tests are carried out using the means and methods described in PCT International Application No. PCT/US97/01609, filed January 29, 1997 which is hereby incorporated by reference.

In these experiments patient-derived segment(s) corresponding to the HIV protease and reverse transcriptase coding regions were either patient-derived segments amplified by the reverse transcription-polymerase chain reaction method (RT-PCR) using viral RNA isolated from viral particles present in the serum of HIV-infected individuals or were mutants of wild type HIV-1 made by site directed mutagenesis of a parental clone of resistance test vector DNA. Isolation of viral RNA was performed using standard procedures (e.g. RNagents Total RNA Isolation System, Promega, Madison WI or RNAzol, Tel-Test, Friendswood, TX). The RT-PCR protocol was divided into two steps. A retroviral reverse transcriptase [e.g. Moloney MuLV reverse transcriptase (Roche Molecular Systems, Inc., Branchburg, NJ), or avian myeloblastosis virus (AMV) reverse transcriptase, (Boehringer Mannheim, Indianapolis, IN)] was used to copy viral RNA into cDNA. The cDNA was then amplified using a thermostable DNA polymerase [e.g. Taq (Roche Molecular Systems, Inc., Branchburg, NJ), Tth (Roche Molecular Systems, Inc., Branchburg, NJ), PrimeZyme (isolated from *Thermus brockianus*, Biometra, Gottingen, Germany)] or a combination of thermostable polymerases as described for the performance of "long PCR" (Barnes, W.M., (1994) Proc. Natl. Acad. Sci, USA 91, 2216-2220) [e.g. Expand High Fidelity PCR System (Taq + Pwo), (Boehringer Mannheim, Indianapolis, IN) OR GeneAmp XL PCR kit (Tth + Vent), (Roche Molecular Systems, Inc., Branchburg, NJ)].

35

The primers, ApaI primer (PDSApa) and AgeI primer (PDSAge) used to amplify the "test" patient-derived segments contained sequences resulting in ApaI and AgeI recognition sites being introduced into the 5' and 3' termini of the PCR product, respectively as described in PCT International Application No. PCT/US97/01609, filed January 29, 1997.

Resistance test vectors incorporating the "test" patient-derived segments were constructed as described in PCT International Application No. PCT/US97/01609, filed January 29, 1997 using an amplified DNA product of 1.5 kB prepared by RT-PCR using viral RNA as a template and oligonucleotides PDSApa (1) and PDSAge (2) as primers, followed by digestion with ApaI and AgeI or the isoschizimer PINAI. To ensure that the plasmid DNA corresponding to the resultant resistance test vector comprises a representative sample of the HIV viral quasi-species present in the serum of a given patient, many (>100) independent E. coli transformants obtained in the construction of a given resistance test vector were pooled and used for the preparation of plasmid DNA.

A packaging expression vector encoding an amphotrophic MuLV 4070A env gene product enables production in a resistance test vector host cell of resistance test vector viral particles which can efficiently infect human target cells. Resistance test vectors encoding all HIV genes with the exception of env were used to transfect a packaging host cell (once transfected the host cell is referred to as a resistance test vector host cell). The packaging expression vector which encodes the amphotrophic MuLV 4070A env gene product is used with the resistance test vector to enable production in the resistance test vector host cell of infectious pseudotyped resistance test vector viral particles.

Resistance tests performed with resistance test vectors were carried out using packaging host and target host cells consisting of the human embryonic kidney cell line 293 (Cell Culture Facility, UC San Francisco, SF, CA) or the Jurkat
5 leukemic T-cell line (Arthur Weiss, UC San Francisco, SF, CA).

Resistance tests were carried out with resistance test vectors using two host cell types. Resistance test vector
10 viral particles were produced by a first host cell (the resistance test vector host cell) that was prepared by transfecting a packaging host cell with the resistance test vector and the packaging expression vector. The resistance
15 test vector viral particles were then used to infect a second host cell (the target host cell) in which the expression of the indicator gene is measured.

The resistance test vectors containing a functional luciferase gene cassette were constructed and host cells
20 were transfected with the resistance test vector DNA. The resistant test vectors contained patient-derived reverse transcriptase and protease sequences that were either susceptible or resistant to the antiretroviral agents, such as nucleoside reverse transcriptase inhibitors, non-
25 nucleoside reverse transcriptase inhibitors and protease inhibitors. The resistance test vector viral particles produced by transfecting the resistance test vector DNA into host cells, either in the presence or absence of protease inhibitors, were used to infect target host cells grown
30 either in the absence of NRTI or NNRTI or in the presence of increasing concentrations of the drug. The amount of luciferase activity produced in infected target host cells in the presence of drug was compared to the amount of luciferase produced in infected target host cells in the
35 absence of drug. Drug resistance was measured as the amount

of drug required to inhibit by 50% the luciferase activity detected in the absence of drug (inhibitory concentration 50%, IC₅₀). The IC₅₀ values were determined by plotting percent drug inhibition vs. log₁₀ drug concentration.

5

Host cells were seeded in 10-cm-diameter dishes and were transfected several days after plating with resistance test vector plasmid DNA and the envelope expression vector. Transfections were performed using a calcium-phosphate precipitation procedure. The cell culture media containing the DNA precipitate was replaced with fresh medium, from one to 24 hours, after transfection. Cell culture media containing resistance test vector viral particles was harvested one to four days after transfection and was passed through a 0.45-µm filter before being stored at -80°C. HIV capsid protein (p24) levels in the harvested cell culture media were determined by an EIA method as described by the manufacturer (SIAC; Frederick, MD). Before infection, target cells (293 and 293/T) were plated in cell culture media. Control infections were performed using cell culture media from mock transfections (no DNA) or transfections containing the resistance test vector plasmid DNA without the envelope expression plasmid. One to three or more days after infection the media was removed and cell lysis buffer (Promega) was added to each well. Cell lysates were assayed for luciferase activity (Fig.3). The inhibitory effect of the drug was determined using the following equation:

$$\% \text{ luciferase inhibition} = 1 - (\text{RLU}_{\text{luc}} [\text{drug}] \div \text{RLU}_{\text{luc}}) \times 100$$

where RLU_{luc} [drug] is the relative light unit of luciferase activity in infected cells in the presence of drug and RLU_{luc} is the Relative Light Unit of luciferase activity in infected cells in the absence of drug. IC₅₀ values were obtained from the sigmoidal curves that were generated from the data by plotting the percent inhibition of luciferase

activity vs. the log₁₀ drug concentration. The drug inhibition curves are shown in (Fig.3).

EXAMPLE 2: Correlating Phenotypic Susceptibility And
5 Genotypic Analysis

Phenotypic susceptibility analysis of patient HIV samples
Resistance test vectors are constructed as described in example 1. Resistance test vectors, or clones derived from the resistance test vector pools, are tested in a phenotypic
10 assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs may comprise members of the classes known as nucleoside-analog reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase
15 inhibitors (NNRTIs), and protease inhibitors (PRIs). The panel of drugs can be expanded as new drugs or new drug targets become available. An IC₅₀ is determined for each resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested is
20 examined and compared to known patterns of susceptibility. A patient sample can be further examined for genotypic changes correlated with the pattern of susceptibility observed.

25 Genotypic analysis of patient HIV samples

Resistance test vector DNAs, either pools or clones, are analyzed by any of the genotyping methods described in Example 2. In one embodiment of the invention, patient HIV sample sequences are determined using viral RNA
30 purification, RT/PCR and ABI chain terminator automated sequencing. The sequence that is determined is compared to control sequences present in the database or is compared to a sample from the patient prior to initiation of therapy, if available. The genotype is examined for sequences that are

different from the control or pre-treatment sequence and correlated to the observed phenotype.

Phenotypic susceptibility analysis of site directed mutants

5 Genotypic changes that are observed to correlate with changes in phenotypic patterns of drug susceptibility are evaluated by construction of resistance test vectors containing the specific mutation on a defined, wild-type (drug susceptible) genetic background. Mutations may be

10 incorporated alone and/or in combination with other known drug resistance mutations that are thought to modulate the susceptibility of HIV to a certain drug or class of drugs. Mutations are introduced into the resistance test vector through any of the widely known methods for site-directed

15 mutagenesis. In one embodiment of this invention the megaprimer PCR method for site-directed mutagenesis is used. A resistance test vector containing the specific mutation or group of mutations is then tested using the phenotypic susceptibility assay described above and the susceptibility

20 profile is compared to that of a genetically defined wild-type (drug susceptible) resistance test vector which lacks the specific mutations. Observed changes in the pattern of phenotypic susceptibility to the antiretroviral drugs tested is attributed to the specific mutations introduced into the

25 resistance test vector.

EXAMPLE 3

Correlating Phenotypic Susceptibility And Genotypic Analysis: P225H

30 Phenotypic analysis of Patient 97-302

A resistance test vector was constructed as described in example 1 from a patient sample designated as 97-302. This patient had been treated with d4T, indinavir and DMP-266 for a period of approximately 10 months. Isolation of viral RNA

35 and RT/PCR was used to generate a patient derived segment

that comprised viral sequences coding for all of PR and aa
1 - 313 of RT. The patient derived segment was inserted
into a indicator gene viral vector to generate a resistance
test vector designated RTV-302. RTV-302 was tested using a
5 phenotypic susceptibility assay to determine accurately and
quantitatively the level of susceptibility to a panel of
anti-retroviral drugs. This panel of anti-retroviral drugs
comprised members of the classes known as NRTIs (AZT, 3TC,
d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), and
10 PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An
IC50 was determined for each drug tested. Susceptibility of
the patient virus to each drug was examined and compared to
known patterns of susceptibility. A pattern of
susceptibility to the NNRTIs was observed for patient sample
15 RTV-302 in which there was significant decrease in
nevirapine susceptibility (increased resistance) and modest
decrease in delavirdine susceptibility (See Figure 8A).
Patient sample 97-302 was examined further for genotypic
changes associated with the observed pattern of
20 susceptibility.

Determination of genotype of patient 97-302

RTV-302 DNA was analyzed by ABI chain terminator automated
sequencing. The nucleotide sequence was compared to the
25 consensus sequence of a wild type clade B HIV-1 (HIV
Sequence Database Los Alamos, NM). The nucleotide sequence
was examined for sequences that are different from the
control sequence. RT mutations were noted at positions
K103N, I135M, T200A, and P225H. K103N is associated with
30 resistance to the NNRTIs and has been shown using the
phenotypic susceptibility assay to be associated with
reduced susceptibility to both delavirdine and nevirapine
to an equal extent. The mutations at I135M and T200A are
known polymorphisms of the wild-type (drug-sensitive)
35 variants of HIV. The mutation, P225H, was characterized

using site directed mutagenesis and phenotypic susceptibility testing to correlate the changes at amino acid 225 with changes in NNRTI phenotypic susceptibility.

5 Site directed mutagenesis

Resistance test vectors were constructed containing the P225H mutation alone and in combination with other known drug resistance mutations (K103N, Y181C) known to modulate the HIV susceptibility to NNRTIs. Mutations were introduced
10 into the resistance test vector using the mega-primer PCR method for site-directed mutagenesis. (Sakar G and Sommar SS (1994) *Biotechniques* 8(4), 404-407). A resistance test vector containing the P225H mutation (P225H-RTV) was tested using the phenotypic susceptibility assay described above
15 and the results were compared to that of a genetically defined resistance test vector that was wild type at position 225. The pattern of phenotypic susceptibility to the NNRTI, delavirdine in the P225H-RTV was altered as compared to wild type. In the context of an otherwise wild
20 type background (i.e. P225H mutation alone) the P225H-RTV was more susceptible to delavirdine than the wild type control RTV. No significant change in nevirapine susceptibility was observed in the P225H-RTV. The P225H mutation was also introduced into a RTV containing
25 additional mutations at K103N, Y181C or both (K103N+Y181C). In all cases, RTVs were more susceptible to inhibition by delavirdine if the P225H mutation was present as compared to the corresponding RTV lacking the P225H mutation (Fig. 8D). In all cases the P225H mutation did not significantly change
30 nevirapine susceptibility (Fig. 8D).

EXAMPLE 4**Correlating Phenotypic Susceptibility And Genotypic Analysis: P236L****Phenotypic analysis of HIV patient 97-268**

5 A resistance test vector was constructed as described in Example 1 from a patient sample designated 97-268. This patient had been treated with AZT and 3TC (NRTIs), indinavir and saquinavir (PRIs) and delavirdine (an NNRTI) for periods varying from 1 month to 2 years. Isolation of viral
10 RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and amino acids 1 - 313 of RT. The patient derived segment was inserted into a indicator gene viral vector to generate a resistance test vector designated RTV-268. RTV-268 was
15 then tested using the phenotypic susceptibility assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs
20 (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for each drug tested. Susceptibility of the patient virus to each drug was examined and compared to the susceptibility of a reference virus. A pattern of
25 susceptibility to the NNRTIs was observed for the patient sample RTV-268 in which the virus sample was observed to be resistant to delavirdine with no resistance to delavirdine. The sample was examined further for genotypic changes associated with the pattern of susceptibility.

30

Genotype of HIV patient 97-268

RTV-268 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of wild type clade B HIV-1. The
35 nucleotide sequence was evaluated for sequences different

from the control sequence. RT mutations were noted at positions M41L, D67N, M184V, T200A, E203D, L210W, T215Y, K219Q, and P236L compared to the control sequence. The mutations at T200A and E203D are known polymorphisms in wild-type (drug-sensitive) variants of HIV. Mutations at positions M41L, D67N, L210W, T215Y, and K219Q are associated with AZT resistance. The mutation at M184V is associated with 3TC resistance. The mutation at P236L is associated with resistance to delavirdine and increased susceptibility to nevirapine (Dueweke et al., Ibid.). In contrast to previous reports, the RTV-268 sample showed no change in nevirapine susceptibility. The mutation, P236L, was characterized using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate changes at amino acid 236 with changes in phenotypic susceptibility.

Site directed mutagenesis

Resistance test vectors were constructed containing the P236L mutation alone and in combination with other known drug resistance mutations (K103N, Y181C) that are known to modulate the susceptibility of HIV-1 to NNRTIs. Mutations were introduced into the resistance test vector using the mega-primer PCR method for site-directed mutagenesis (Sakar and Sommar, Ibid.). A resistance test vector containing the P236L mutation (P236L-RTV) was tested using the phenotypic susceptibility assay and the results were compared to that of a genetically defined resistance test vector that was wild type at position 236. P236L-RTV exhibited changes in NNRTI phenotypic susceptibility. In the context of an otherwise wild type background (i.e. P236L mutation alone) the P236L-RTV is less susceptible to delavirdine than a wild type reference RTV. In contrast to Dueweke et al. no significant change in nevirapine susceptibility was observed for P236L-RTV. The P236L mutation was also introduced into a RTV containing mutations at K103N, Y181C or both

(K103N+Y181C). In all cases, the RTV's were less susceptible (more resistant) to delavirdine if the P236L mutation was present as compared to the corresponding RTV lacking the P236L mutation. In all cases the P236L mutation
5 did not significantly alter nevirapine susceptibility.

Example 5

Correlating Phenotypic Susceptibility And Genotypic Analysis: G190S

10 Phenotypic analysis of HIV patient 97-644

A resistance test vector was constructed as described in Example 1 from a patient sample designated 97-644. This patient had been treated with d4T (NRTI), indinavir (PRI) and efavirenz (NNRTI) for a period varying from 5 to 17
15 months. Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and amino acids 1 - 313 of RT. The patient derived segment was inserted into a indicator gene viral vector to generate a resistance test
20 vector designated RTV-644. RTV-644 was then tested using the phenotypic susceptibility assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT,
25 3TC, d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for each drug tested. Susceptibility of the patient virus to each drug was examined and compared to the susceptibility of a
30 reference virus. A pattern of susceptibility to the NNRTIs was observed for the patient sample RTV-644 in which the virus sample was observed to be resistant to nevirapine with little or no resistance to delavirdine. The sample was examined further for genotypic changes associated with
35 the pattern of susceptibility.

Genotype of HIV patient 97-644

RTV-644 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of wild type clade B HIV-1. The
5 nucleotide sequence was evaluated for sequences different from the control sequence. RT mutations were noted at positions K101E and G190S compared to the control sequence. The mutations at T200A and E203D are known polymorphisms in wild-type (drug-sensitive) variants of
10 HIV. The mutation at K101E is associated with resistance to some but not all NNRTIs. The mutation, G190A but not specifically G190S is associated with nevirapine and loviride resistance. The mutations G190S and G190A were characterized using site directed mutagenesis and in vitro
15 phenotypic susceptibility testing to correlate changes at amino acid 190 with changes in phenotypic susceptibility.

Site directed mutagenesis

Resistance test vectors were constructed containing the
20 G190S and G190A mutations. Mutations were introduced into the resistance test vector using the mega-primer PCR method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the G190S or G190A mutations (G190S-RTV, or G190A-RTV) were tested using the
25 phenotypic susceptibility assay and the results were compared to that of a genetically defined resistance test vector that was wild type at position G190. G190S-RTV and G190A-RTV exhibited changes in NNRTI phenotypic susceptibility. In the context of an otherwise wild type
30 background these RTVs were markedly less susceptible to nevirapine and slightly more susceptible to delavirdine than a wild type reference RTV.

Example 6**Predicting Response to Non-nucleoside Reverse Transcriptase Inhibitors by Characterization of Amino Acid Changes in HIV-1 Reverse Transcriptase****5 Phenotypic and genotypic correlation of mutations at amino acid 236 of HIV-1 Reverse Transcriptase**

In one embodiment of this invention, changes in the amino acid at position 236 of the reverse transcriptase protein of HIV-1 is evaluated using the following method comprising:

- 10 (i) collecting a biological sample from an HIV-1 infected subject; (ii) evaluating whether the biological sample contains nucleic acid encoding HIV-1 reverse transcriptase having a mutation at codon 236. The presence of a mutation at codon 236 (P236L) is correlated with a reduction in
15 delavirdine susceptibility and little or no change in nevirapine susceptibility.

The biological sample comprises whole blood, blood components including peripheral mononuclear cells (PBMC),
20 serum, plasma (prepared using various anticoagulants such as EDTA, acid citrate-dextrose, heparin), tissue biopsies, cerebral spinal fluid (CSF), or other cell, tissue or body fluids. In another embodiment, the HIV-1 nucleic acid (genomic RNA) or reverse transcriptase protein can be
25 isolated directly from the biological sample or after purification of virus particles from the biological sample. Evaluating whether the amino acid at position 236 of the HIV-1 reverse transcriptase is mutated, can be performed using various methods, such as direct characterization of
30 the viral nucleic acid encoding reverse transcriptase or direct characterization of the reverse transcriptase protein itself. Defining the amino acid at position 236 of reverse transcriptase can be performed by direct characterization of the reverse transcriptase protein by conventional or novel
35 amino acid sequencing methodologies, epitope recognition by

antibodies or other specific binding proteins or compounds. Alternatively, the amino acid at position 236 of the HIV-1 reverse transcriptase protein can be defined by characterizing amplified copies of HIV-1 nucleic acid encoding the reverse transcriptase protein. Amplification of the HIV-1 nucleic acid can be performed using a variety of methodologies including reverse transcription-polymerase chain reaction (RT-PCR), NASBA, SDA, RCR, or 3SR as would be known to the ordinarily skilled artisan. Evaluating whether the nucleic acid encoding HIV reverse transcriptase has a mutation at codon 236 can be performed by direct nucleic acid sequencing using various primer extension-chain termination (Sanger, ABI/PE and Visible Genetics) or chain cleavage (Maxam and Gilbert) methodologies or more recently developed sequencing methods such as matrix assisted laser desorption-ionization time of flight (MALDI-TOF) or mass spectrometry (Sequenom, Gene Trace Systems). Alternatively, the nucleic acid sequence encoding amino acid position 236 can be evaluated using a variety of probe hybridization methodologies, such as genechip hybridization sequencing (Affymetrix), line probe assay (LiPA; Murex), and differential hybridization (Chiron).

In a preferred embodiment of this invention, evaluation of whether amino acid position 236 of HIV-1 reverse transcriptase was wild type or mutant was carried out using a phenotypic susceptibility assay using resistance test vector DNA prepared from the biological sample. In one embodiment, plasma sample was collected, viral RNA was purified and an RT-PCR methodology was used to amplify a patient derived segment encoding the HIV-1 protease and reverse transcriptase regions. The amplified patient derived segments were then incorporated, via DNA ligation and bacterial transformation, into an indicator gene viral vector thereby generating a resistance test vector.

Resistance test vector DNA was isolated from the bacterial culture and the phenotypic susceptibility assay was carried out as described in Example 1. The results of the phenotypic susceptibility assay with a patient sample having a P236L mutation. The nucleic acid (DNA) sequence of the patient derived HIV-1 protease and reverse transcriptase regions from patient sample 268 was determined using a fluorescence detection chain termination cycle sequencing methodology (ABI/PE). The method was used to determine a consensus nucleic acid sequence representing the combination of sequences of the mixture of HIV-1 variants existing in the subject sample (representing the quasispecies), and to determine the nucleic acid sequences of individual variants.

Phenotypic susceptibility profiles of patient samples and site directed mutants showed that delavirdine and nevirapine susceptibility correlated with the absence of RT mutations at positions 103, 181 or 236 of HIV-1 reverse transcriptase. Phenotypic susceptibility profiles of patient samples and site directed mutants showed a significant reduction in delavirdine susceptibility (increased resistance) and little or no reduction in nevirapine susceptibility correlated with a mutation in the nucleic acid sequence encoding the amino acid leucine (L) at position 236 of HIV-1 reverse transcriptase and the absence of mutations at positions 103 and 181.

Phenotypic susceptibility profiles of patient samples and site directed mutants showed no additional reduction in delavirdine or nevirapine susceptibility (increased resistance) with the amino acid proline at position 236 when the RT mutations at positions 103, 181 or 103 and 181 were present (K103N, Y181C, or K103N + Y181C). However, phenotypic susceptibility profiles of patient samples and site directed mutants showed an additional reduction in

delavirdine susceptibility (increased resistance) and little or no additional reduction in nevirapine susceptibility with the amino acid leucine (L) at position 236 in addition to the RT mutations associated with NNRTI resistance (K103N, Y181C, or K103N + Y181C).

Phenotypic and genotypic correlation of mutations at amino acid 225 of HIV-1 Reverse Transcriptase

Phenotypic susceptibility profiles of patient samples and site directed mutants showed no change in susceptibility to delavirdine or nevirapine when the amino acid proline (P) was present at position 225 of HIV-1 reverse transcriptase in the absence of RT mutations associated with NNRTI resistance (K103N, Y181C). However, phenotypic susceptibility profiles of patient samples and site directed mutants showed an increase in delavirdine susceptibility and little or no change nevirapine susceptibility when the amino acid histidine (H) was present at position 225 in the absence of RT mutations (K103N, Y181C) associated with NNRTI resistance.

Phenotypic susceptibility profiles of patient samples and site directed mutants showed no additional reduction in delavirdine susceptibility or nevirapine susceptibility when the amino acid proline (P) at position 225 was present in addition to the RT mutations associated with NNRTI resistance (K103N, Y181C, or K103N + Y181C). In contrast phenotypic susceptibility profiles of patient samples and site directed mutants showed an increase in delavirdine susceptibility and little or no change in nevirapine susceptibility when the amino acid histidine (H) was present at position 225 in the presence of RT mutations associated with NNRTI resistance (K103N, Y181C, or K103N + Y181C).

Phenotypic and genotypic correlation of mutations at amino acid 190 of HIV-1 Reverse Transcriptase

Phenotypic susceptibility profiles of patient samples and site directed mutants showed no change in susceptibility to
5 delavirdine or nevirapine when the amino acid glycine (G) at position 190 was present in the absence of RT mutations associated with NNRTI resistance (K103N, Y181C). Phenotypic susceptibility profiles of site directed mutants showed an increase in delavirdine susceptibility and a decrease in
10 nevirapine susceptibility when the amino acid alanine (A) was present at position 190 in the absence of RT mutations associated with NNRTI resistance. Phenotypic susceptibility profiles of patient samples and site directed mutants showed an increase in delavirdine susceptibility and a decrease in
15 nevirapine susceptibility when the amino acid serine (S) was present at position 190 in the absence of RT mutations associated with NNRTI resistance.

EXAMPLE 8

- 20 **Using Resistance Test Vectors and Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility And Resistance in HIV: Y181I**
Preparation of resistant test vectors and phenotypic analysis of patient 98-964 HIV samples
- 25 A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-964. This patient had been previously treated with ddI, d4T, AZT, 3TC, ddC, (NRTIs), saquinavir and nelfinavir (PRIs) and nevirapine (an NNRTI) and HU. Isolation of viral RNA and
30 RT/PCR was used to generate a patient derived segment that comprised viral sequence coding for all of PR and aa 1- 313 of RT. The PDS was inserted into an indicator gene viral vector to generate a resistance test vector designated RTV-964. RTV-964 was then tested in a phenotypic assay to
35 determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the

classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-964 in which there was a moderate decrease (10_fold) in delavirdine susceptibility and a significant decrease (750-fold) in nevirapine susceptibility.

Determination of genotype of patient HIV samples
RTV-964 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions M41L, K43E, D67N, K70R, L74I, V75S, Y181I, R211T, T215Y, D218E, and K219Q compared to the control sequence. M41L, D67N, K70R, L74I, V75S, T215Y, and K219Q are associated with NRTI resistance. A mutation at R211T is a known polymorphism in the sequence among different wild-type (drug-sensitive) variants of HIV. Y181I had previously been shown to be associated with high level resistance to nevirapine. We examined the mutation, Y181I, using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to anti-retroviral drugs in HIV

The Y181I mutation was introduced into the resistance test vector using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid). A resistance test vector containing the Y181I mutation (Y181I -RTV) was then

tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at position 181. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in the Y181I-RTV. On a wild type background (i.e. Y181I mutation alone) the Y181I-RTV displayed a moderate loss of susceptibility (20-fold) to delavirdine and a significant loss of susceptibility (740-fold) to nevirapine compared to a wild type control RTV. The Y181I- RTV showed wild-type susceptibility (1.4-fold) to efavirenz.

EXAMPLE 9

- 15 Using Resistance Test Vectors And Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility And Resistance in HIV: Y188

Preparation of resistant test vectors and phenotypic analysis of patient 97-300 HIV samples

- A resistance test vector was constructed as described in Example 1 from a patient sample designated 97-300. This patient had been previously treated with d4T and 3TC (NRTIs), indinavir (a PRI) and efavirenz (an NNRTI).
- 25 Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1 - 313 of RT. The PDS was inserted into an indicator gene viral vector to generate a resistance test vector designated RTV-300. RTV-300 was then
- 30 tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delavirdine, efavirenz and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for the resistance test
- 35 vector pool for each drug tested. The pattern of susceptibility to all of the drug tested was examined and

compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-300 in which there was moderate decrease (25-fold) in delavirdine susceptibility and a substantial decrease
5 (greater than 800-fold) in nevirapine susceptibility.

Determination of genotype of patient HIV samples
RTV-300 DNA analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the
10 consensus sequence of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NH). The genotype was examined for sequence that are different from the control sequence. Mutations were noted at positions K32N, M184V and Y188L compared to the control sequence. The mutation at
15 M184V is associated with 3TC resistance. Y188L had previously been shown to be associated with high level resistance to efavirenz. Other mutations at position Y188 (i.e Y188C and Y188H) have been reported to have been selected for by treatment with several NNRTIs (E-ePseU, E-
20 EPS, HEPT, Nevirapine, BHAP, U-8720E, TIBO R82913, Loviride). We examined the mutation, Y188L, using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with
phenotype.

25 Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV

The Y188L mutation was introduced into the resistance test
30 vector using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.): A resistance test vector containing the Y188L mutation (Y188L-RTV) was then tested using the phenotypic assay described earlier and the results were compared to those determined using a
35 genetically defined resistance test vector that was wild type at position 188. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in the Y188L-RTV. On a wild type

background (i.e. Y188L mutation alone) the Y188L-RTV displayed a slight loss of susceptibility (9-fold) to delavirdine and substantial loss of susceptibility (greater than 800-fold) to nevirapine and a significant loss of susceptibility (109-fold) to efavirenz compared to a wild type control RTV. The approximate 100-fold loss of susceptibility to efavirenz was not as high as had been previously reported.

- 10 Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV

The Y188C mutation was introduced into the resistance test vector using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). A resistance test vector containing the Y188C mutation (Y188C-RTV) was then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at position 188. We determined the pattern of phenotypic susceptibility to the NNRTIs., delavirdine, nevirapine and efavirenz, in the Y188C-RTV. On a wild type background (i.e. Y188C mutation alone) the Y188C-RTV displayed a slight loss of susceptibility (3-fold) to delavirdine and a moderate loss of susceptibility (30-fold) to nevirapine and efavirenz (20-fold) compared to a wild type control RTV.

- 30 Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV

The Y188H mutation was introduced into the resistance test vector using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). A resistance test vector containing the Y188H mutation (Y188H-RTV) was then tested using the phenotypic assay described earlier and the results were compared to those determined using a

genetically defined resistance test vector that was wild type at position 188. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine and nevirapine, in the Y188H-RTV. On a wild type background
5 (i.e. Y188H mutation alone) the Y188H-RTV displayed a moderate loss of susceptibility (3.5-fold) to nevirapine compared to a wild type control RTV. The phenotypic susceptibility of Y188H to efavirenz was not determined.

10 **EXAMPLE 10**

Using Resistance Test Vectors And Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility And Resistance in HIV: E138 and Y188

15 **Preparation of resistant test vectors and phenotypic analysis of patient 97-209 HIV samples**

A resistance test vector was constructed as described in Example 1 from a patient sample designated 97-209. This patient had been previously treated with AZT, ddI, d4T and
20 3TC (NRTIs), indinavir (a PRIs) and adefovir. Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1 - 313 of RT. The PDS was inserted into an indicator gene viral vector to generate resistance test
25 vector designated RTV-209. RTV-209 was then tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC),
30 NNRTIs (delavirdine, efavirenz and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC₅₀ was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drug tested. The pattern of susceptibility to all of the
35 drugs tested was examined and compared to known patterns of

susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-209 in which there was a moderate decrease (75-fold) in delavirdine susceptibility and a substantial decrease (greater than 800-fold) in
5 nevirapine susceptibility.

Determination of genotype of patient HIV samples

RTV-209 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the
10 consensus sequence of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions A62V, S68G, V76I, F77L, F116Y, E138A, Q151M, M184V, Y188L and E291D
15 compared to the control sequence. The mutations at A62V, V75I, F77L, F116Y, Q151M and M184V are associated with NRTI resistance. A mutation at E138K had previously been shown to be associated with resistance to several NNRTIs and a mutation at Y188L had previously been shown to be associated
20 with a decrease in susceptibility to efavirenz. We examined the mutations Y188L and E138A using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

25

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV

The E138A mutation alone and in combination with Y188L was
30 introduced into resistance test vectors using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the E138A mutation (E138A-RTV) or the E138 mutation along with the Y188L mutation (E138A-Y188L-RTV) were then tested using
35 the phenotypic assay described earlier and the results were

compared to those determined using a genetically defined resistance test vector that was wild type at positions 188 and 138. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in the E138A-RTV, Y188L-RTV and E138-Y188L-RTV. On a wild type background (i.e. E138A mutation alone) the E138A-RTV displayed wild-type susceptibility to delavirdine (1.6-fold), nevirapine (1.3-fold) and efavirenz (1.4-fold). The Y188L-RTV displayed a slight loss of susceptibility (greater than 800-fold) to nevirapine and a significant loss of susceptibility (110-fold) to efavirenz. The E138A-Y188L-RTV displayed a moderate loss of susceptibility (75-fold) to delavirdine and efavirenz (88-fold) and a substantial loss of susceptibility to nevirapine (greater than 800-fold) compared to a wild type control RTV. The combination of mutations resulted in an increased effect on delavirdine susceptibility compared to the effect observed for each mutation alone.

20 **EXAMPLE 11**

Using Resistance Test Vectors And SiteDirected Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility And Resistance in HIV: A98

25 **Preparation of resistant test vectors and phenotypic analysis of patient 98-675 HIV samples**

A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-675. This patient had been previously treated with ddI, AZT, and 3TC (NRTIs), and saquinavir and nelfinavir (PRIs). Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1 - 313 of RT. The PDS was inserted into an indicator gene viral vector to generate a resistance test vector designated RTV-675. RTV-675 was then tested in a

phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI
5 and ddC), NNRTIs (delavirdine, efavirenz and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all
10 of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-675 in which wild-type susceptibility (2.1-fold) was observed for delavirdine and a slight decrease (6-fold) in nevirapine susceptibility was observed.

15

Determination of genotype of patient HIV samples

RTV-675 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV
20 Sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions M41L, S48t, L74V, A98G, M184V and T215Y are associated with NRTI resistance. A mutation at A98G had previously been shown to
25 be associated with resistance to nevirapine. We examined the mutation A98G using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

30 **Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV**

The A98G mutation into the resistance test vector using the
35 mega-primer method for site-directed mutagenesis (Sakar and

Sommar, Ibid.). A resistance test vector containing the A98G mutation (A98G-RTV) was then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at position 98. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in the A98G-RTV. On a wild type background (i.e. A98G mutation alone) the A98G RTV displayed a slight loss of susceptibility to delavirdine (3-fold), nevirapine (8-fold) and efavirenz (3-fold) compared to a wild type control RTV.

Example 12 Using Resistance Test Vectors and Site Directed Mutants to Correlate Genotypes And Phenotypes Associated with NNRTI Drug Susceptibility and Resistance in HIV: A98 and G190

Preparation of resistant test vectors and phenotypic analysis of patient B HIV samples.

A resistant test vector was constructed as described in Example 1 from a patient sample designated B. The anti-retroviral treatment this patient received is unknown. Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1-313 of RT. The PDS was inserted into an indicator gene viral vector to generate a resistant test vector designated RTV-B. Individual clones of the RTV-B pool were selected and then tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC₅₀ was determined for the resistance test vector clone for each

drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-B clone 1 in which there was an
5 increase in susceptibility (0.55-fold) to delaviridine, a substantial loss of susceptibility (640-fold) to nevirapine and significant loss of susceptibility (250-fold) to efavirenz.

10 **Determination of genotype of patient HIV samples**

RTV-B clone 1 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was
15 examined for sequences that are different from the control sequence. Mutations were noted at positions M41L, A98G, M184V, L210W, R211?, T215Y, E297A and G190S compared to the control sequence. M41L, M184V, L210W and T215Y are associated with NRTI resistance. A mutation at A98G had
20 previously been shown to be associated with resistance to nevirapine. A mutation at position G190A had previously been shown to be associated with changes in susceptibility to nevirapine. Other changes at position 190 (i.e. E, Q, and T) have also been reported. We examined the mutations
25 A98G and G190S, using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to anti-viral drugs in HIV

The A98 and G190S mutations were introduced alone or in
5 combination into the resistance test vector using the megaprimer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the A98G mutation (A98G-RTV), the G190S mutation (G190S-RTV) and both mutations (A98G-G190S-RTV) were then tested using the
10 phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at position 98 and 190. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in the
15 three vectors. On a wild type background (i.e. A98G mutation alone) the A98G-RTV displayed a slight loss of susceptibility to delavirdine (3-fold), nevirapine (8-fold) and efavirenz (3-fold) compared to a wild type control RTV. On a wild type background (i.e. G190S mutation alone) the
20 G190S-RTV displayed increased susceptibility (0.5-fold) to delavirdine, a moderate loss of susceptibility (75-fold) to nevirapine and a slight loss of susceptibility (8-fold) to efavirenz compared to a wild type control RTV. The A98G-G190S-RTV displayed increased susceptibility (0.8-fold) to
25 delavirdine, but a substantial loss of susceptibility to both nevirapine (greater than 800-fold) and efavirenz (greater than 250-fold) compared to a wild type control RTV. Although only a slight loss of susceptibility to efavirenz was observed for the individual mutations, the combination
30 of A98G and G190S resulted in a substantial loss of susceptibility to efavirenz. Likewise, this combination of mutation resulted in a greater loss of susceptibility to nevirapine than the sum of the two mutations alone.

EXAMPLE 13

**Using Resistance Test Vectors and Site Directed Mutants
Correlate Genotypes And Phenotypes Associated With NNRTI
Drug Susceptibility And Resistance in HIV: Y181 and A98**

5

**Preparation of resistant test vectors and phenotypic
analysis of patient 98-1057 samples**

A resistance test vector was constructed as described in
Example 1 from a patient sample designated 98-1057. This
10 patient had been previously treated with ddI, d4T, AZT, and
3TC (NRTIs), saquinavir and indinavir (PRIs) and delavirdine
(an NNRTI). Isolation of viral RNA and RT/PCR was used to
generate a patient derived segment that comprised viral
sequences coding for all of PR and aa 1-313 RT. The PDS was
15 inserted into an indicator gene viral vector to generate
resistance test vector designated RTV-1057. RTV-1057 was
then tested in a phenotypic assay to determine accurately
and quantitatively the level of susceptibility to a panel of
anti-retroviral drugs. This panel of anti-retroviral drugs
20 comprised members of the classes known as NRTIs (AZT, 3TC,
d4T, ddI, and ddC), NNRTIs (delavirdine, efavirenz and
nevirapine) and PRIs (indinavir, nelfinavir, ritonavir, and
saquinavir). An IC₅₀ was determined for the resistance test
vector pool for each drug tested. The pattern of
25 susceptibility to all of the drugs tested was examined and
compared to known patterns of susceptibility. A pattern of
susceptibility to the NNRTIs was observed for patient RTV-
1057 in which there was a moderate decrease in delavirdine
(35-fold) susceptibility and a significant decrease (610-
30 fold) in nevirapine susceptibility.

Determination of genotype of patient HIV samples

RTV-1057 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV
5 Sequence Database, Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions T39A, M41L, A62V, D67E, T69SST, A98G, I135T, Y181C, T200I and T215Y
10 compared to the control sequence M41L, A62V, D67E, T69SST, and T215Y are associated with NRTI resistance. Mutations at positions I135T and T200I are known polymorphisms in the sequence among different wild-type (drug-sensitive) variants of HIV. Y181C and A98G have been previously shown to be associated with resistance to certain NNRTIs. We examined
15 the mutations Y181C and A98G using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

20 **Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to anti-retroviral drugs in HIV**

The Y181C and A98G mutations were introduced alone and in combination into resistance test vectors using the megaprimer method for site-directed mutagenesis (Sakar and
25 Sommar, Ibid.). Resistance test vectors containing the Y181C mutation (Y181C-RTV) and the A98G mutation (A98G-RTV) and both mutations (Y181C-A98G-RTV) were then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined
30 resistance test vector that was wild type at position 181 and 98. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in the three vectors. On a wild type background (i.e. Y181C mutation alone) the Y181C-RTV displayed moderate
35 loss of susceptibility (35-fold) to delavirdine, a significant loss of susceptibility (161-fold) to nevirapine

and a slight loss of susceptibility (3-fold) to efavirenz compared to a wild type control RTV. The A98G-RTV displayed a slight loss of susceptibility to delavirdine (3-fold), nevirapine (8-fold) and efavirenz (3-fold) compared to a wild type control RTV. The Y181C-A98G-RTV displayed significant loss of susceptibility (240-fold) to delavirdine, a substantial loss of susceptibility (greater than 800-fold) to nevirapine and a slight loss of susceptibility (7-fold) to efavirenz compared to a wild type control RTV. These data indicated that the combination of the two mutations, Y181C and A98G, resulted in a greater loss of susceptibility to both delavirdine and nevirapine than the sum of effects observed for these two mutations individually.

15

EXAMPLE 14

Using Resistant Test Vectors and Site Directed Mutants to Correlate Genotypes and Phenotypes Associated with NNRTI Drug Susceptibility and Resistance in HIV: K101 and G190

20

Preparation of resistant test vectors and phenotypic analysis of patients 98-644 and 98-1060 HIV samples

A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-644. This patient had been previously treated with d4T (an NNRTI), indinavir (a PRI and efavirenz (an NNRTI). A second resistance test vector was constructed as described in Example 1 from a patient sample designated 98-1060. This patient had been previously treated with d4T (an NNRTI). indinavir (a PRI) and efavirnez (an NNRTI). Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1-313 of RT. The PDS was inserted into an indicator gene viral vector to generate resistance test vectors designated RTV-644 and RTV-1060. RTV-644 and RTV-1060 were then tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs

comprised members of the classes known as NNRTIs (AZT, 3TC, d4T, ddI, and ddC), NNRTIs (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-644 in which there was a very slight (2.5-fold) decrease in delavirdine susceptibility and a significant (600-fold) decrease in nevirapine susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-644 in which there was a very slight (2.5-fold) decrease in delavirdine susceptibility and a significant (600-fold) decrease in nevirapine susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-1060 in which wild-type susceptibility (1.5-fold) to delavirdine was observed. A significant decrease in efavirenz susceptibility (900-fold) and a substantial decrease to nevirapine (greater than 800-fold) susceptibility was observed for RTV-1060.

Determination of genotype of patient HIV samples

RTV-644 and RTV-1060 DNA were analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions K101E and G190S for RTV-644 compared to the control sequence and mutations were noted at positions K101E, G190S, T200A and T215Y for RTV-1060 compared to the control sequence. The sequence at position T215 was a mixture of wild-type and mutation. A mutation at position K101E had been previously shown to be associated with resistance to several NNRTIs including high level resistance to delavirdine. A mutation at position G190A had previously been shown to be associated with changes in susceptibility to nevirapine. Other changes at position 190 (i.e. E, Q and T) have also been reported. We

examined the mutations K101E and G190S, using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

5

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV

10 The K101E and G190S mutations were introduced alone and in combination into resistance test vectors using the megaprimer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the K101E mutation (K101E-RTV), the G190S mutation (G190S-RTV) were then tested using the phenotypic assay described
15 earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at positions 101 and 190. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in all three vectors.
20 On a wild type background (i.e. K101E mutation alone) the K101E-RTV displayed a slight loss of susceptibility (5-fold) to delavirdine and efavirenz (5-fold) and a moderate loss of susceptibility (12-fold) to nevirapine compared to a wild type control RTV. The K101E-G190S-RTV displayed increased
25 susceptibility to delavirdine (0.5-fold), a moderate loss of susceptibility to nevirapine (75-fold) and a slight loss of susceptibility (7.6-fold) to efavirenz compared to a wild type control RTV. The K101E-G190S-RTV displayed wild-type susceptibility (1.4-fold) to delavirdine and a substantial
30 loss of susceptibility to both nevirapine (greater than 800-fold) and efavirenz (greater than 250-fold) compared to a wild type control RTV.

In this example, the combination of mutations, G190S and
35 K101E, displayed a novel phenotypic pattern. The combination resulted in the reversal of the effect on delavirdine susceptibility observed for the G190S mutation

alone and a greater than additive effect on the susceptibility for both nevirapine and efavirenz.

EXAMPLE 15

- 5 Using Resistance Test Vectors And Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug susceptibility And Resistance in HIV: V108I

10 Preparation of resistant test vectors and phenotypic analysis of patient 98-652 HIV samples

- A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-652. This patient had no previous anti-retroviral treatment. Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1 - 313 or RT. The PDS was inserted into an indicator gene viral vector to generate a resistance test vector designated RTV-652. RTV-652 was then tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir and saquinavir). An IC₅₀ was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-652 in which increase susceptibility (0.97-fold) to delavirdine was observed and a slight decrease (5-fold) in nevirapine susceptibility was observed.

Determination of genotype of patient HIV samples

- 35 RTV-652 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was

examined for sequences that are different from the control sequence. Mutations were noted at positions M41L, V108I, I135T, L210W, R211K and T215D compared to the control sequence. M41L, L210W and T215D are associated with NRTI resistance. Mutations at positions I135T and R211K are known polymorphisms in the sequence among different wild-type (drug-sensitive) variants of HIV. V108I is known to be associated with resistance to several NNRTIs. We examined the mutation V108I using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to antiretroviral drugs in HIV

The V108I mutation was introduced into the resistance test vector using the mega-primer method for site directed mutagenesis (Sakar and Sommar, Ibid.). A resistance test vector containing the V108I mutation (V108I-RTV) was then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at position 108. We determined the pattern of phenotypic susceptibility to the NNRTIs, delaviridine, nevirapine and efavirenz, in the V108I -RTV. On a wild type background (i.e. V108I mutation alone) the V108I -RTV displayed wild-type susceptibility (1.3-fold) to delaviridine and efavirenz (1.7-fold) and a slight loss of susceptibility (3-fold) to nevirapine compared to a type control RTV.

EXAMPLE 16

Using Resistance Test Vectors And Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility And Resistance in HIV: K103 and K101 and G190

Preparation of resistant test vectors and phenotypic analysis of patient 98-955 HIV samples

A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-955. This patient had been previously treated with nelfinavir (a PRI). Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of PR and aa 1 - 313 of RT. The PDS was inserted into an indicator gene viral vector to generate a resistance test vectors designated RTV-955. RTV-955 was then tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delaviridine, efavirenz and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-955 in which there was a slight decrease (4-fold) in delaviridine susceptibility and a significant decrease (530-fold) in nevirapine susceptibility.

25

Determination of genotype of patient HIV samples

RTC-955 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions K20R, V35I, A62V, D67N, T69D, V75I, F77L, K101E, K103N, Y115F, F116Y, Q151M, I167V, Y181C, M184V, G190A, I202V, R211K, F214L, T215V, and K219Q compared to the control sequence. Mutations at positions K101E, K103N, Y181C, G190A, and F214 L were mixtures of wild-type and the mutation. A62V, D67N, T69D, V75I, F77L, Y115F, F116Y, Q151M, M184V, T215V and K219Q are associated

with NRTI resistance. Mutations at V35I, R211K and F214L are known polymorphism in the sequence among different wild-type (drug sensitive) variants of HIV. a mutation at position K101E had been previously shown to be associated with resistance to the NNRTIs. A mutation at Y181I had previously been shown to be associated with high level resistance to nevirapine. a mutation at K103N had previously been shown to be associated with resistance to the three NNRTIs, delaviridine and nevirapine and efavirenz. We examined the mutations K101E, J103N and G190A using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to anti-retroviral drugs in HIV

The K101E, K103N and G190A mutations were introduced alone and in combination into resistance test vectors using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the K101E mutation (K101E-RTV), the K103N mutation (K0103N-RTV), the G190 mutation (g190A-RTV and two mutations (K101E-G190A-RTV) and (K103N-G190A-RTV) were then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at positions 101, 103 and 190. We determined the pattern of phenotypic susceptibility to the NNRTIs, delaviridine, nevirapine, and efavirenz, in all 5 vectors. On a wild type background (i.e. K101E mutation alone) the K101E-RTV displayed a slight loss (5-fold) os susceptibility to delavirdine and efavirenz (5-fold) and a moderate loss of susceptibility (12-fold) to nevirapine (55-fold) and efavirenz(30-fold) compared to a wild type control RTV. On a wild type background (i.e. G190A mutation alone) the G190A -RTV displayed increased susceptibility (8-fold) efavirenz compared to a wild type control RTV. The K101E-G190A-RTV displayed wild-type susceptibility (2-fold) to

delavirdine, substantial loss of susceptibility (greater than 800-fold) to nevirapine and a significant loss of susceptibility (120-fold) to efavirenz compared to a wild type control RTV. The K103N-G190-RTV displayed a moderate
5 loss of susceptibility (40-fold) to delavirdine, substantial loss of susceptibility (greater than 800-fold) to nevirapine and a significant loss of susceptibility (215-fold) to efavirenz compared to a wild type control RTV. The introduction of a second mutation to a vector containing the
10 G190A resulted in the reversal of the effect on delavirdine susceptibility observed for the G190A mutation alone. The G190-a mutation displayed an increased susceptibility to delviridine. However, the addition of either K10E or K103N to the G190A mutation resulted in a slight loss of
15 susceptibility to delavirdine. Furthermore, the combination of G190A and K101E resulted in a greater than additive effect on the loss of susceptibility to nevirapine and efavirenz. Lastly, these data indicated that the combination of the two mutations G190A and K103N resulted in a greater
20 loss of susceptibility to both nevirapine and efavirenz than the sum of effects observed for these two mutations individually.

EXAMPLE 17

25 Using Test Vectors And Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility An Resistance in HIV: V106 and V189 and V181 and F227

30 Preparation of resistant test vectors and phenotypic analysis of patient 98-1033 and 98-757 HIV samples

A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-1033. This patient had been previously treated with AZT, d5T, 3TC and
35 ddI (NRTI), saquinavir, indinavir and nelfinavir (PRIs and nevirapine (an NNRTI). a second resistance test vector was constructed as described in Example 1 from a sample obtained from the same patient at a different time point and

designated 98-757. This patient had received an additional 8 weeks of treatment with nevirapine 9an NNRTI) d4T (an NRTI), and saquinavir and nelfinavir (PRIs). Isolation of viral RNA and RT/PCR was used to generate a patient derived
5 segment that comprised viral sequences coding for all of PR and aa 1 - 313 of RT. The PDS was inserted into an indicator gene viral vector to generate resistance test vectors designated RTV-1033 and RTV-757. RTV-1033 and RTV-757 were then tested in a phenotypic assay to determine
10 accurately and quantitatively the level of susceptibility to a panel of anti-retroviral drugs. This panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), and PRIs (indinavir, nelfinavir, ritonavir, and
15 saquinavir). An IC50 was determined for the resistance test vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for patient RTV-
20 1033 in which there was a moderate decrease (30-fold) in delavirdine susceptibility and a substantial decrease (greater than 800-fold) in nevirapine susceptibility and a significant decrease (200-fold) in efavirenz susceptibility. A pattern of susceptibility to the NNRTIs was observed for
25 patient RTV-757 in which there was a slight decrease (10-fold) in delavirdine susceptibility and a substantial decrease (greater than 800-fold) in nevirapine susceptibility.

Determination of Genotype of Patient HIV Samples

RTV-1033 and RTV-757 DNA were analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV Sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence. Mutations were noted at positions V35I, D67N, T69D, K70R, V106A, V189L, T200A, I202T, R211K, T215F, D218E, K219Q, H221Y, F227L, L228H and R284 for RTV-1033 compared to the control sequence. Mutations were noted at positions V35I, D67N, T69D, K70R, V106A, V108I, L109V, Y108C, V189L, T200A, I202T, R211K, T215F, D218E, K219Q, H221Y, L228H, L283I and R284K for RTV-757 compared to the control sequence. The sequences at positions V106A, V108I and L109V were a mixture of wild-type and mutation. D67N, T69D, K70R, T215F and K219Q are associated with NRTI resistance. Mutations at V35I, T200A, R211K and R284K are known polymorphisms in the sequence among different wild-type (drug-sensitive) variants of HIV. A mutation at V106A had previously been shown to be associated with increase resistance to nevirapine. A mutation at V189I had previously been shown to be associated with NNRTI resistance but a mutation to L at this position had not been previously reported to be associated with NNRTI resistance. A mutation at V108I had previously been shown to be associated with increased resistance to both delavirdine and nevirapine. A mutation at Y181C had also previously been shown to be associated with increased resistance to both delavirdine and nevirapine. We examined the mutations V106A, V189L, V181C and F227L using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to anti-retroviral drugs in HIV

The mutations V106A, V189L, V181C and F227L were introduced alone and in combination into resistance test vectors using the mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the V106A mutation (V106A-RTV), the V189L mutation (V189L-RTV), the V181C mutation (V181C-RTV) and F227L mutation (F227L-RTV) and two mutations (V106A-V181C-RTV) and (V106A-V189L-RTV) and (V106A-F227-RTV) and (V181C-F227-RTV) and three mutations, (V106A-V181C-F227L-RTV) were then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at positions 106, 189, 181 and 227. We determined the pattern of phenotypic susceptibility to the NNRTIs, delavirdine, nevirapine and efavirenz, in all nine vectors. On a wild type background (i.e. V106A mutation alone) the V106A-RTV displayed a slight loss (5-fold) of susceptibility to delavirdine and a moderate loss of susceptibility (60-fold) to nevirapine and wild-type susceptibility (1.7-fold) to efavirenz compared to a wild type control RTV. On a wild type background (i.e. V189L mutation alone) the V189-RTV displayed wild type susceptibility to delavirdine (1.8-fold), nevirapine (1.3-fold) and efavirenz (1.3-fold) compared to a wild type control RTV. On a wild type background (i.e. V181C mutation alone) the V181C-RTV displayed a significant loss of susceptibility (100-fold) to delavirdine and a substantial loss of susceptibility (greater than 800-fold) to nevirapine and a slight loss of susceptibility (4-fold) to efavirenz compared to a wild type control RTV. On a wild type background (i.e. F227L mutation alone) the F227L-RTV displayed increased susceptibility (0.03-fold) to delavirdine and efavirenz (0.48-fold) and a slight loss of

susceptibility (3-fold) to nevirapine compared to a wild type control RTV. The V106A-Y181C-RTV displayed a significant loss of susceptibility (100-fold) to delavirdine, a substantial loss of susceptibility (greater than 800-fold) to nevirapine and slight loss of susceptibility (4-fold) to efavirenz compared to a wild type control RTV. The V106A-V189L-RTV displayed a slight loss of susceptibility (3-fold) to delavirdine, a moderate loss of susceptibility (50-fold) to nevirapine and wild-type susceptibility (1-fold) to efavirenz compared to a wild type control RTV. The V106A-F227-RTV displayed a slight loss of susceptibility (3-fold) to delavirdine, a substantial loss of susceptibility (greater than 800-fold) to nevirapine and a slight loss of susceptibility (8-fold) to efavirenz compared to a wild type control RTV. The Y181C-F227L-RTV displayed increased susceptibility (0.89-fold) to delavirdine and efavirenz (0.99-fold) and a significant loss of susceptibility (285-fold) to nevirapine compared to a wild type control RTV. The V106A-Y181C-F227L-RTV displayed a moderate loss (50-fold) of susceptibility to delavirdine and a substantial loss of susceptibility (greater than 800-fold) to nevirapine and a slight loss of susceptibility (12-fold) to efavirenz compared to a wild type control RTV.

25 **EXAMPLE 18**

Using Resistance Test Vectors And Site Directed Mutants To Correlate Genotypes And Phenotypes Associated With NNRTI Drug Susceptibility And Resistance In HIV: Y188 and L100 and K103

30

Preparation of resistance test vectors and phenotypic analysis of patient 98-1058 HIV samples

A resistance test vector was constructed as described in Example 1 from a patient sample designated 98-1058. This patient had been previously treated with ddI, d4T, AZT, 3TC,

35

ddC and abacavir (NRTIs), indinavir and amprenavir (PRIs) and nevirapine (an NNRTI). Isolation of viral RNA and RT/PCR was used to generate a patient derived segment that comprised viral sequences coding for all of RP and aa 1 -
5 313 of RT. The PDS was inserted into an indicator gene viral vector to generate a resistance test vector designated RTV-1058. Individual clones of RTV-1058 were selected and were then tested in a phenotypic assay to determine accurately and quantitatively the level of susceptibility to
10 a panel of anti-retroviral drugs. The panel of anti-retroviral drugs comprised members of the classes known as NRTIs (AZT, 3TC, d4T, ddI and ddC), NNRTIs (delavirdine and nevirapine), an PRIs (indinavir, nelfinavir, ritonavir, and saquinavir). An IC50 was determined for the resistance test
15 vector pool for each drug tested. The pattern of susceptibility to all of the drugs tested was examined and compared to known patterns of susceptibility. A pattern of susceptibility to the NNRTIs was observed for clones 4, 5 and 10 from patient RTV-1058. Clone 4 displayed a
20 significant loss of susceptibility (85-fold) for delavirdine and a substantial loss of susceptibility (greater than 800-fold) for nevirapine. Clone 5 displayed a substantial loss of susceptibility (250-fold) to delavirdine and a significant loss of susceptibility (120-fold) to nevirapine.
25 Clone 10 displayed a substantial loss of susceptibility (greater than 250-fold) to delavirdine and (greater than 800-fold) to nevirapine.

Determination of genotype of patient HIV samples

30 RTV-1058 DNA was analyzed by ABI chain terminator automated sequencing. The nucleotide sequence was compared to the consensus sequence of a wild type clade B HIV-1 (HIV sequence Database Los Alamos, NM). The genotype was examined for sequences that are different from the control sequence.
35 Mutations were noted at positions M41L, A62V, D67N, T69SST,

L74V, L100I, K103N, V118I, I135T, T200S, L210W, R211K and T215Y compared to the control sequence. L74V and L100I were mixtures of wild-type and mutation. Clone 4 contained mutations at positions K103N and Y188L. Clone 5 contained 5 mutations at positions L100I and K103N. Clone 10 contained mutations at positions L100I, K103N and Y188L. M41L, A62V, D67N, T69SST, L74V, L210W and T215Y are associated with NRTI resistance. Mutations at positions I135T, T200S and R211T are known polymorphisms in the sequence among different 10 wild-type (drug-sensitive) variants of HIV. A mutation at L100I had previously been shown to be associated with resistance to delavirdine and nevirapine. A mutation at K103N had previously been shown to be associated with resistance to delavirdine, nevirapine and efavirenz. We 15 examined the mutations, Y188L, L100I and K103N, using site directed mutagenesis and in vitro phenotypic susceptibility testing to correlate the observed changes in genotype with phenotype.

20 **Site directed mutagenesis is used to confirm the role of specific mutations in phenotypic susceptibility to anti-retroviral drugs in HIV**

The mutations Y188L, L100I and K103N were introduced alone and in combination into resistance test vectors using the 25 mega-primer method for site-directed mutagenesis (Sakar and Sommar, Ibid.). Resistance test vectors containing the Y188L mutation (Y188L-RTV), the L100I mutation (L100I-RTV), the K103N mutation (K103N-RTV), the two mutations (K103N-Y188L-RTV) and (L100I-K103N-RTV), and the three mutations 30 (L100I-K103N-Y188L-RTV) were then tested using the phenotypic assay described earlier and the results were compared to those determined using a genetically defined resistance test vector that was wild type at positions 188, 100, and 103. We determined the pattern of phenotypic 35 susceptibility to the NNRTIs, delavirdine, nevirapine and

efavirenz, in all 6 vectors. On a wild type background (i.e. Y188L mutation alone) the Y188L-RTV displayed a slight loss of susceptibility (9-fold) to delavirdine, a substantial loss of susceptibility (greater than 800-fold) to nevirapine and a moderate loss of susceptibility (110-fold) to efavirenz compared to a wild type control RTV. On a wild type background (i.e. L100I mutation alone) the L100I-RTV displayed a moderate loss of susceptibility (30-fold) to delavirdine and efavirenz (10-fold) and a slight loss of susceptibility (10-fold) and a slight loss of susceptibility (3-fold) to nevirapine compared to a wild type control RTV. On a wild type background (i.e. K103M mutation alone) the K103N-RTV displayed moderate loss of to delavirdine susceptibility (50-fold), nevirapine (55-fold) and efavirenz (30-fold) compared to a wild type control RTV. The K103N-Y188L-RTV displayed substantial loss of susceptibility to delavirdine (greater than 250-fold), nevirapine (greater than 800-fold) and efavirenz (greater than 250-fold) compared to a wild type control RTV. The L100I-K103N-RTV displayed substantial loss of susceptibility (greater than 250-fold) to delavirdine and efavirenz (greater than 250-fold) and a moderate loss of susceptibility (70-fold) to nevirapine compared to a wild type control RTV. The L100I-K103N-Y188L-RTV displayed substantial loss of susceptibility to delavirdine (greater than 250-fold), nevirapine (greater than 800-fold), and efavirenz (greater than 250-fold) compared to a wild type control RTV. Novel combinations resulted in unpredeicted resistance patterns than were different from those patterns observed for the each mutation alone.

What is claimed is:

1. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy
5 of an HIV-infected patient comprising:
 - (a) collecting a plasma sample from the HIV-infected patient; and
 - (b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase
10 having a mutation at codon 236;
in which the presence of the mutation correlates with decreased susceptibility to delavirdine and little or no change in nevirapine susceptibility.
- 15 2. The method of claim 1, wherein the mutation at codon 236 codes for a leucine.
3. The method of claim 1, wherein reverse transcriptase has an additional mutation(s) at codon 103, codon 181
20 or a combination thereof.
4. The method of claim 3, wherein the mutation at codon 103 encodes an asparagine (N) and the mutation at codon 181 encodes a cysteine ©).
25
5. The method of claim 1, wherein the HIV-infected patient is being treated with an antiretroviral agent.
6. A method of assessing the effectiveness of
30 antiretroviral therapy of an HIV-infected patient comprising:
 - (a) collecting a biological sample from an HIV-infected patient; and

- (b) evaluating whether the biological sample comprises nucleic acid encoding HIV reverse transcriptase having a mutation at codon 225;
in which the presence of the mutation correlates with
5 an increase in delavirdine susceptibility and little or no change in nevirapine susceptibility.
7. The method of claim 6, wherein the mutated codon 225 encodes a histidine (H).
- 10 8. The method of claim 6, wherein the HIV-infected patient is being treated with an antiretroviral agent.
- 15 9. The method of claim 6, wherein the reverse transcriptase has an additional mutation(s) at codon 103, 181 or a combination thereof.
- 20 10. A method of assessing the effectiveness of antiretroviral therapy of an HIV-infected patient comprising:
(a) collecting a biological sample from an HIV-infected patient; and
(b) evaluating whether the biological sample comprises nucleic acid encoding HIV reverse
25 transcriptase having a mutation at codon 190;
in which the presence of the mutation correlates with an increase in delavirdine susceptibility and a decrease in nevirapine susceptibility.
- 30 11. The method of claim 10, wherein the mutation at codon 190 encodes an alanine or a serine.
- 35 12. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:

- (a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 236 and a mutation at codon 103 and/or 181 and an indicator gene into a host cell;
- 5 (b) culturing the host cell from step (a);
- (c) measuring the indicator in a target host cell; and
- (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in
- 10 the absence of the candidate antiretroviral drug compound;

wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

15

13. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:

- (a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 225 and a mutation at codon 103 and an indicator into a host cell;
- 20 (b) culturing the host cell from step (a);
- (c) measuring the indicator in a target host cell; and
- 25 (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;

30 wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c) ; or at step (c)

14. A method for assessing the biological effectiveness of
a candidate HIV antiretroviral drug compound
comprising:

- (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 236 and an indicator gene into a host cell;
- 5 (b) culturing the host cell from step (a);
- (c) measuring expression of the indicator gene in a target host cell; and
- (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are
- 10 carried out in the absence of the candidate anti-viral drug compound,
- wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at
- 15 steps (b) - (c); or at step (c).

15. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:
- 20 (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 225, and an indicator gene into a host cell;
- (b) culturing the host cell from step (a);
- 25 (c) measuring expression of the indicator gene in a target host cell; and

- (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound,
- 5 wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).
- 10 16. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:
- 15 (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 190, and an indicator gene into a host cell;
- (b) culturing the host cell from step (a);
- (c) measuring expression of the indicator gene in a target host cell; and
- 20 (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound,
- 25 wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

17. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 190 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
18. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 225 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
19. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 236 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
20. A resistance test vector of claim 17, wherein the patient-derived segment having a mutation at codon 190 further comprises a mutation at codon 103.
21. A resistance test vector of claim 18, wherein the patient-derived segment having a mutation at codon 225 further comprises a mutation at codon 103.
22. The resistance test vector of claim 17, wherein the patient-derived segment having a mutation at codon 236 further comprises a mutation at codon 103 and/or 181.
23. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy of an HIV-infected comprising:

- (a) collecting a plasma sample from an HIV-infected patient; and
(b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase having a mutation at codon 230;
in which the presence of the mutation correlates with decreased susceptibility to delavirdine and nevirapine.
24. The method of claim 23, wherein the mutation at codon 230 codes for a leucine (L).
25. The method of claim 23, wherein reverse transcriptase has an additional mutation(s) at codon 181 or a combination thereof.
26. The method of claim 25, wherein the mutation at codon 281 encodes a cysteine (C).
27. The method of claim 23, wherein the HIV-infected patient is being treated with an antiretroviral agent.
28. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy of an HIV-infected comprising:
(a) collecting a plasma sample from an HIV-infected patient; and
(b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase having a mutation at codon 181;
in which the presence of the mutation correlates with decreased susceptibility to delavirdine and nevirapine and little or no change in efavirenz susceptibility.
29. The method of claim 28, wherein the mutation at codon 181 codes for a cysteine (c).

30. The method of claim 28, wherein reverse transcriptase has an additional mutation(s) at codon 98, codon 106, codon 227 or a combination thereof.
- 5 31. The method of claim 30, wherein the mutation at codon 98 encodes a glycine (G), the mutation at codon 106 encodes an alanine (A) and the mutation at codon 227 encodes a leucine (L)
- 10 32. The method of claim 28, wherein the HIV-infected patient is being treated with an antiretroviral agent.
33. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy of an HIV-infected comprising:
- 15 (a) collecting a plasma sample from an HIV-infected patient; and
- (b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase
- 20 having a mutation at codon 188;
- in which the presence of the mutation correlates with decreased susceptibility to delavirdine and nevirapine and efavirenz.
- 25 34. The method of claim 33, wherein the mutation at codon 188 codes for a leucine (L), cysteine (C) or histidine (H).
35. The method of claim 33, wherein reverse transcriptase has an additional mutation(s) at codon 138, codon 103, codon 100 or a combination thereof.
- 30 36. The method of claim 35, wherein the mutation at codon 138 encodes an alanine (A), the mutation at codon 103

encodes an asparagine (N) and the mutation at codon 100 encodes an isoleucine (I).

37. The method of claim 33, wherein the HIV-infected patient is being treated with an antiretroviral agent.
38. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy of an HIV-infected comprising:
- (a) collecting a plasma sample from an HIV-infected patient; and
- (b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase having a mutation at codon 190;
- in which the presence of the mutation correlates with increased susceptibility to delavirdine and decreased susceptibility to nevirapine and efavirenz.
39. The method of claim 38, wherein the mutation at codon 190 codes for an alanine (A) or a serine (S).
40. The method of claim 38, wherein reverse transcriptase has an additional mutation(s) at codon 98, codon 101 or codon 103 or a combination thereof.
41. The method of claim 40, wherein the mutation at codon 98 encodes a glycine (G), 101 encodes a glutamic acid (E) and 103 encodes an asparagine (N).
42. The method of claim 38, wherein the HIV-infected patient is being treated with an antiretroviral agent.

43. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy of an HIV-infected comprising:
- 5 (a) collecting a plasma sample from an HIV-infected patient; and
- (b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase having a mutation at codon 106;
- 10 in which the presence of the mutation correlates with decreased susceptibility to delavirdine and nevirapine and little or no change in efavirenz susceptibility.
44. The method of claim 43, wherein the mutation at codon 106 encodes an alanine (A).
- 15 45. The method of claim 43, wherein reverse transcriptase has an additional mutation(s) at codon 227 and codon 189 or a combination thereof.
- 20 46. The method of claim 45, wherein the mutation at codon 227 encodes a leucine (L) and 189 encodes a leucine (L).
47. The method of claim 43, wherein the HIV-infected patient is being treated with an antiretroviral agent.
- 25 48. A method of assessing the effectiveness of non-nucleoside reverse transcriptase antiretroviral therapy of an HIV-infected comprising:
- 30 (a) collecting a plasma sample from an HIV-infected patient; and
- (b) evaluating whether the plasma sample contains nucleic acid encoding HIV reverse transcriptase having a mutation at codon 103;

in which the presence of the mutation correlates with decreased susceptibility to delavirdine and nevirapine and efavirenz.

- 5 49. The method of claim 48, wherein the mutation at codon 103 codes for asparagine (N).
- 10 50. The method of claim 48, wherein reverse transcriptase has an additional mutation(s) at codon 100 or a combination thereof.
51. The method of claim 50, wherein the mutation at codon 100 encodes an isoleucine (I).
- 15 52. The method of claim 48, wherein the HIV-infected patient is being treated with an antiretroviral agent.
53. A method for assessing the biological effectiveness of candidate HIV antiretroviral drug compound comprising:
- 20 (a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 230 and a mutation at codon 181 and an indicator gene into a host cell;
- (b) culturing the host cell from step (a);
- 25 (c) measuring the indicator in a target host cell; and
- (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;
- 30 wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

54. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:
- 5 (a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 181 and a mutation at codon 98 and/or 106 and/or 227 and an indicator gene into a host cell;
 - (b) culturing the host cell from step (a);
 - 10 (c) measuring the indicator in a target host cell; and
 - (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;
 - 15 wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).
- 20 55. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:
- (a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 188 and a mutation at codon 138 and/or 103 and/or 100 and an indicator gene into a host cell;
 - 25 (b) culturing the host cell from step (a);
 - (c) measuring the indicator in a target host cell; and
 - 30 (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;

wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c) or at step (c).

- 5 56. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:

(a) introducing a resistance test vector comprising a patient-derived segment further comprising a
10 mutation

at codon 190 and a mutation at codon 98 and/or 101 and/or 103 and an indicator gene into a host cell;

(b) culturing the host cell from step (a);

(c) measuring the indicator in a target host cell; and

5 (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;

10 wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

57. A method for assessing the biological effectiveness of
15 a candidate HIV antiretroviral drug compound comprising:

(a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 106 and a mutation at codon 227
20 and/or 189 and an indicator gene into a host cell;

(b) culturing the host cell from step (a);

(c) measuring the indicator in a target host cell; and

(d) comparing the measurement of the indicator from
25 step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;

wherein a test concentration of the candidate
30 antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

58. A method for assessing the biological effectiveness of
a candidate HIV antiretroviral drug compound
35 comprising:

- (a) introducing a resistance test vector comprising a patient-derived segment further comprising a mutation at codon 103 and a mutation at codon 100 and an indicator gene into a host cell;
- 5 (b) culturing the host cell from step (a);
- (c) measuring the indicator in a target host cell; and
- (d) comparing the measurement of the indicator from step (c) with the measurement of the indicator measured when steps (a) - (c) are carried out in the absence of the candidate antiretroviral drug compound;
- 10

wherein a test concentration of the candidate antiretroviral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

15

59. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:

- (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 230 and an indicator gene into a host cell;
- 20 (b) culturing the host cell from step (a);
- (c) measuring expression of the indicator gene in a target host cell; and
- 25 (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound,
- 30

wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

60. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:
- 5 (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 227 and an indicator gene into a host cell;
 - (b) culturing the host cell from step (a);
 - 10 (c) measuring expression of the indicator gene in a target host cell; and
 - (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound,
 - 15 wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) -(c); or at step (c).
- 20 61. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:
- 25 (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 188 and an indicator gene into a host cell;
 - (b) culturing the host cell from step (a);
 - (c) measuring expression of the indicator gene in a target host cell; and
 - 30 (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound,

wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

5 62. A method for assessing the biological effectiveness of a candidate HIV antiretroviral drug compound comprising:

10 (a) introducing a resistance test vector comprising a patient-derived segment which encodes the reverse transcriptase having a mutation at codon 189 and an indicator gene into a host cell;

(b) culturing the host cell from step (a);

(c) measuring expression of the indicator gene in a target host cell; and

15 (d) comparing the expression of the indicator gene from step (c) with the expression of the indicator gene measured when steps (a) - (c) are carried out in the absence of the candidate anti-viral drug compound,

20 wherein a test concentration of the candidate anti-viral drug compound is present at steps (a) - (c); at steps (b) - (c); or at step (c).

25 63. A resistant test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 230 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.

30 64. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 227 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.

35

65. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 188 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
66. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutation at codon 189 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
67. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutations at codon 181, codon 98, codon 106 and codon 227 or a combination thereof and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
68. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutations at codon 106, codon 227, and codon 189 or a combination thereof and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.
69. A resistance test vector comprising an HIV patient-derived segment further comprising reverse transcriptase having a mutations at codon 103 and codon 101 and an indicator gene, wherein the expression of the indicator gene is dependent upon the patient derived segment.

117


70. The resistance test vector of claim 17, wherein the patient-derived segment having a mutation at codon 190 further comprises mutations at codon 98 and codon 101 or a combination thereof.

5

71. The resistance test vector of claim 63, wherein the patient-derived segment having a mutation at codon 230 further comprises a mutation at codon 181.

10 72. The resistance test vector of claim 65, wherein the patient-derived segment having a mutation at codon 188 further comprises mutations at codon 138, codon 103 and codon 190 or a combination thereof.

Resistance Test Vector



Resistance Test Vector




FIGURE 1

Two Cell Assay




FIGURE 2

Drug Resistance: NRTI, NNRTI, PI




FIGURE 3

Drug Susceptibility Test Profile: Patient 487




FIGURE 4

Drug Susceptibility Test Profile: Site Directed Mutants




FIGURE 5

Drug Susceptibility Test Profile: Patient 268




FIGURE 6

Drug Susceptibility Test Profile: Site Directed Mutants




FIGURE 7

Drug Susceptibility Test Profile: Patient 302




FIGURE 8A

Drug Susceptibility Test Profile: Patient 780




FIGURE 8B

Drug Susceptibility Test Profile: Patient 302 Virus Clones




FIGURE 8C

Drug Susceptibility Test Profile: Site Directed Mutants




FIGURE 8D

Drug Susceptibility Test Profile: Patient 644




FIGURE 9A




FIGURE 9B

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/11629**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/5, 6, 8, 91.1, 91.2, 91.4, 91.42, 91.5, 91.51, 320.1, 455, 456, 457, 465; 536/23.1, 23.2, 23.72

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE CLERCQ, E. Development of resistance of human immunodeficiency virus (HIV) to anti-HIV agents: how to prevent the problem. International Journal of Antimicrobial Agents. June 1997, Vol. 9, pages 21-36, especially pages 26-28.	1-5, 10-11, 28-44, 47-52
Y		6-9, 12-14, 16-17, 19-22, 45-46, 54-58, 61, 65, 67-70, 72
Y	PELEMANS et al. Characteristics of the Pro225His mutation in human immunodeficiency virus type 1 (HIV-1) reverse transcriptase that appears under selective pressure of dose-escalating quinoxaline treatment of HIV-1. J. Virology. November 1997, Vol. 71, No. 11, pages 8195-8203, especially pages 8195, 8197, 8201, Table 2.	6-9, 13, 15, 18, 21

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

25 AUGUST 1999

Date of mailing of the international search report

30 SEP 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DIANA JOHANNSEN

Telephone No. (703) 308-0196

JOYCE BRIDGERS
PARALEGAL SPECIALIST
CHEMICAL MATRIX

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/11629

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BALZARINI et al. A novel mutation (F227L) arises in the reverse transcriptase of human immunodeficiency virus type 1 on dose-escalating treatment of HIV type 1-infected cell cultures with the nonnucleoside reverse transcriptase inhibitor thiocarboxanilide UC-781. AIDS Research And Human Retroviruses. 10 February 1998, Vol. 14, No. 3, pages 255-260, especially Table 1.	45-46, 57, 60, 64
Y	BALZARINI et al. Zidovudine-resistant human immunodeficiency virus type 1 strains subcultured in the presence of both lamivudine and quinoxaline HBY 097 retain marked sensitivity to HBY 097 but not to lamivudine. J. Of Infectious Diseases. November 1997, Vol. 176, pages 1392-1397, especially Table 2.	45-46, 57, 62, 66
Y	SHI et al. A recombinant retroviral system for rapid in vivo analysis of human immunodeficiency virus type I susceptibility to reverse transcriptase inhibitors. Antimicrobial Agents and Chemotherapy. December 1997, Vol. 41, No. 12, pages 2781-2785, especially pages 2781-2783.	12-22, 54-58, 60-62, 64-69, 72
Y	STRAIR et al. Recombinant retroviral systems for the analysis of drug resistant HIV. Nucleic Acids Research. 1993, Vol. 21, No. 20, pages 4836-4842, especially pages 4836-4838, 4840.	12-22, 54-58, 60-62, 64-69, 72

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/11629

A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

C12Q 1/66, 1/68, 1/70; C12P 19/34; C12N 15/00, 15/64, 15/85; C07H 21/04

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

435/5, 6, 8, 91.1, 91.2, 91.4, 91.42, 91.5, 91.51, 320.1, 455, 456, 457, 465; 536/23.1, 23.2, 23.72

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

USPAT, WPI, CAPLUS, AIDSLINE, MEDLINE, BIOSIS, EMBASE, SCISEARCH, LIFESCI
search terms: HIV, reverse transcriptase, mutations, nevirapine, delavirdine, BHAPs, efavirenz, DMP-266

THIS PAGE BLANK (USPTO)